

# GEOMETRIC CONSIDERATION OF DUALITY IN MULTIOBJECTIVE OPTIMIZATION WITH SET FUNCTIONS

JUN YULL LEE

## 1. Introduction.

In ordinary scalar convex optimization, the Lagrange multiplier theorem asserts the existence of a supporting hyperplane for the epigraph of the primal map[5]. On the other hand, in multiobjective optimization, the corresponding theorem [7, Theorem 2.3] implies the existence of a conical variety (i.e., a translation of a cone) which supports the epigraph of primal map. In this paper, we show that a similar assertion is true for the multiobjective programming problem with set functions.

## 2. Multiobjective Programming Problem with Set Functions.

Let  $(X, \mathcal{A}, \mu)$  be a finite, atomless measure space and  $L^1(X, \mathcal{A}, \mu)$  be separable. Then, by considering characteristic function  $\chi_\Omega$  of  $\Omega$  in  $\mathcal{A}$ , we can embed  $\mathcal{A}$  into  $L^\infty(X, \mathcal{A}, \mu)$ . In this setting for  $\Omega, \Lambda \in \mathcal{A}$ , and  $\alpha \in I = [0, 1]$ , there exists a sequence, called a *Morris sequence*,  $\{\Gamma_n\} \subset \mathcal{A}$  such that

$$\chi_{\Gamma_n} \xrightarrow{w^*} \alpha \chi_\Omega + (1 - \alpha) \chi_\Lambda,$$

where  $\xrightarrow{w^*}$  denotes the *weak\**-convergence of elements in  $L^\infty(X, \mathcal{A}, \mu)$  [6].

A subfamily  $\mathcal{S}$  is said to be *convex* if for every  $(\alpha, \Omega, \Lambda) \in I \times \mathcal{S} \times \mathcal{S}$  and every Morris sequence  $\{\Gamma_n\}$  associated with  $(\alpha, \Omega, \Lambda)$  in  $\mathcal{A}$ , there exists a subsequence  $\{\Gamma_{n_k}\}$  of  $\{\Gamma_n\}$  in  $\mathcal{S}$ . In [1], if  $\mathcal{S} \subseteq \mathcal{A}$  is convex, then the *weak\**-closure  $cl(\mathcal{S})$  of  $\chi_{\mathcal{S}}$  in  $L^\infty(X, \mathcal{A}, \mu)$  is the *weak\**-closed convex hull of  $\chi_{\mathcal{S}}$ , and  $\overline{\mathcal{A}} = \{f \in L^\infty : 0 \leq f \leq 1\}$ .

---

Received July 13, 1995.



DEFINITION 2.1. Let  $\mathcal{S}$  be a convex subfamily of  $\mathcal{A}$ . Let  $K$  be a convex cone of  $R^n$ . A set function  $H : \mathcal{S} \rightarrow R^n$  is called  $K$ -convex, if given  $(\alpha, \Omega_1, \Omega_2) \in I \times \mathcal{S} \times \mathcal{S}$  and Morris-sequence  $\{\Gamma_n\}$  in  $\mathcal{A}$  associated with  $(\alpha, \Omega_1, \Omega_2)$ , there exists a subsequence  $\{\Gamma_{n_k}\}$  of  $\{\Gamma_n\}$  in  $\mathcal{S}$  such that

$$\limsup_{k \rightarrow \infty} H(\Gamma_{n_k}) \leq_K \alpha H(\Omega_1) + (1 - \alpha)H(\Omega_2),$$

where  $\limsup$  is taken over each component. And  $x <_K y$  denotes  $y - x \in \text{int}(K)$ ,  $x \leq_K y$  denotes  $y - x \in K \setminus \{0\}$ , and  $x \leq_K y$  denotes  $y - x \in K$ .

DEFINITION 2.2. A set function  $H = (H_1, H_2, \dots, H_n) : \mathcal{S} \rightarrow R^n$  is called weak\*-continuous on  $\mathcal{S}$  if for each  $f \in \text{cl}(\mathcal{S})$  and for each  $j = 1, 2, \dots, n$ , the sequence  $\{H_j(\Omega_k)\}$  converges to the same limit for all  $\{\Omega_k\}$  with  $\chi_{\Omega_k} \xrightarrow{w^*} f$ .

Now multiobjective programming problem with set functions can be described as follows:

$$\begin{aligned} & \text{Min}_D \quad F(\Omega) \\ \text{(P)} \quad & \text{subject to } \Omega \in \mathcal{S} \\ & \text{and } G(\Omega) \leq_Q 0, \end{aligned}$$

which has been defined as the problem finding all feasible efficient  $D$ - or properly efficient  $D$ -solution with respect to the pointed closed convex cones  $D$  and  $Q$  of Euclidean spaces  $R^p$  and  $R^m$  with nonempty interiors,  $D^\circ$  and  $Q^\circ$ , respectively. That is, letting  $\mathcal{S}' = \{\Omega \in \mathcal{S} : G(\Omega) \leq_Q 0\}$ , we want to find  $\Omega^* \in \mathcal{S}'$  such that

$$(F(\mathcal{S}') - F(\Omega^*)) \cap (-D) = \{0\}, \quad \emptyset \text{ if } 0 \notin D$$

or

$$\text{cl}(p(F(\mathcal{S}') + D - F(\Omega^*))) \cap (-D) = \{0\}, \quad \emptyset \text{ if } 0 \notin D,$$

where the set  $p(S) = \{\alpha y : \alpha > 0, y \in S\}$  is the projecting cone for a set  $S \subset R^p$ . We denote the set of efficient  $D$ -solutions by  $\mathcal{E}(F(\mathcal{S}'), D)$  and the set of properly efficient  $D$ -solutions by  $\mathcal{PE}(F(\mathcal{S}'), D)$ .

For the primal problem (P), we assume that  $F : \mathcal{S} \rightarrow R^p$ ,  $G : \mathcal{S} \rightarrow R^m$  are  $D$ -convex,  $Q$ -convex, respectively and weak\*-continuous. Under this assumptions we have the Lagrange multiplier theorem as in usual multiobjective optimization problems. The set of  $p \times m$  matrices  $\{M \in R^{p \times m} : MQ \subset D\}$  is denoted by  $\mathcal{M}$ .



THEOREM 2.3. [3] Let  $\Omega^*$  be a properly efficient  $D$ -solution to the problem (P). If there is  $\Omega_o \in \mathcal{S}$  such that  $G(\Omega_o) <_Q 0$ , then there exists  $M^* \in \mathcal{M}$  such that

- (1)  $F(\Omega^*) \in \text{Min}_D\{F(\Omega) + M^*G(\Omega) : \Omega \in \mathcal{S}\}$
- (2)  $M^*F(\Omega^*) = 0$ .

In fact,  $F(\Omega^*) \in \text{Min}_{Dcl}(F(\Omega) + M^*G(\Omega) : \Omega \in \mathcal{S})$ .

The generalized Slater's constraint qualification in Theorem 2.3 that there exists  $\Omega_o \in \mathcal{S}$  such that  $G(\Omega_o) <_Q 0$  is assumed in the sequel.

The primal problem (P) is embedded into a family of perturbed problems:

$$\begin{aligned} & \text{Min}_D \quad F(\Omega) \\ (P_u) \quad & \text{subject to } \Omega \in \mathcal{S} \\ & \text{and } G(\Omega) \leq_Q u. \end{aligned}$$

We denote by  $\mathcal{S}(u)$  the set  $\{\Omega \in \mathcal{S} : G(\Omega) \leq_Q u\}$ , and by  $Y(u)$  the set  $F(\mathcal{S}(u))$ .

DEFINITION 2.4. Perturbed (or primal) maps are defined on  $R^m$  by

$$\begin{aligned} W(u) &= \text{Min}_D F(\mathcal{S}(u)) \\ \text{and } \overline{W}(u) &= \text{Min}_{Dcl}(F(\mathcal{S}(u))) \end{aligned}$$

The original problem (P) can be therefore regarded as determining  $F^{-1}(W(0)) \cap \mathcal{S}$ . However, more satisfactory results are obtained if  $\overline{W}$  is used instead.

For each  $M \in \mathcal{M}$ , we define certain maps for (P) on  $\mathcal{M}$  by

$$\begin{aligned} \Phi(M) &= \text{Min}_D\{F(\Omega) + MG(\Omega) : \Omega \in \mathcal{S}\} \\ \overline{\Phi}(M) &= \text{Min}_{Dcl}\{F(\Omega) + MG(\Omega) : \Omega \in \mathcal{S}\} \end{aligned}$$

The map  $\Phi$  and  $\overline{\Phi}$  are called dual maps for (P).



REMARK 2.5.

- (1)  $MG(\cdot) : S \rightarrow R^p$  is  $D$ -convex on  $S$ .
- (2)  $L(\cdot, M) = F(\cdot) + MG(\cdot)$  is  $D$ -convex and  $w^*$ -continuous.
- (3)  $cl(\{F(S) + MG(\Omega)\})$  is  $D$ -convex subset of  $R^p$
- (4) For each  $M \in \mathcal{M}$ , we have

$$cl(\{L(\Omega, M) : \Omega \in S\}) + D = \overline{\Phi}(M) + D,$$

since  $cl(\{L(\Omega, M) : \Omega \in S\})$  is compact and  $D$ -convex.

- (5) For any  $u$  with  $S(u) \neq \emptyset$ ,  $[clY(u)] + D = \overline{W}(u) + D$ .

The relationship between the primal map  $\overline{W}$  and the dual map  $\overline{\Phi}$  now can be established.

THEOREM 2.6. [4] For any  $M \in \mathcal{M}$ , the following equalities hold.

$$\overline{\Phi}(M) = \text{Min}_D \bigcup_{u \in \zeta} (\overline{W}(u) + Mu) = \text{Min}_D \bigcup_{u \in \zeta^o} (\overline{W}(u) + Mu)$$

where  $\zeta = \{u \in R^m : S(u) \neq \emptyset\}$  and  $\zeta^o = \{u \in R^m : \{\Omega \in S : G(\Omega) <_Q u\} \neq \emptyset\}$ .

COROLLARY 2.7. If  $\Omega^*$  is a properly efficient  $D$ -solution to the problem (P) with generalized Slater's constraint qualification, then there exists an  $M^* \in \mathcal{M}$  such that

$$F(\Omega^*) \in \overline{\Phi}(M^*) \cap \Phi(M^*) \subset \text{Min}_D [\bigcup_{u \in \zeta} (\overline{W}(u) + M^*u)].$$

*proof.* The proof is an immediate consequence of Theorems 2.3 and 2.6.

### 3. A Supporting Conical Variety.

All assumptions on  $F, G, D$  and  $Q$  of section 1 and 2 are inherited. Furthermore,  $D$  is assumed to be a polyhedral convex cone. Then  $D$  is the set of all solutions to some finite system of homegenous weak linear inequalities. Hence there exists an  $s \times p$  matrix  $U_1$  such that

$$(1) \quad D = \{y \in R^p : U_1 \cdot y \geq 0\},$$

where the row vectors of  $U_1$  are generators of  $D^o$ . Since  $D$  is pointed, the  $s \times p$  matrix  $U_1$  has the full rank  $p$ . Here  $\mathcal{M} = \{M \in R^{m \times p} : MQ \subset D\}$ .



DEFINITION 3.1. The  $D$ -epigraph of  $W$  is defined by

$$D\text{-epi } W = \{(u, y) \in R^m \times R^p : u \in \zeta, y \in W(u) + D\}.$$

Here  $W(u) = \text{Min}_D Y(u)$  and  $\zeta = \{u : S(u) \neq \emptyset\}$  were introduced in section 2. The  $D$ -epigraph of  $\bar{W}$  is defined similarly. Unlike the case of ordinary convex multiobjective optimization problem,  $D\text{-epi } W$  may not be a closed convex set in  $R^m \times R^p$ . But we can see that  $cl(D\text{-epi } W)$  is convex. The following lemma is similar to Lemma 2.4 [7].

LEMMA 3.2. For a given  $M \in \mathcal{M}$ , the following conditions are equivalent :

- (i)  $F(\Omega) + MG(\Omega) \not\leq_D F(\Omega^*) + MG(\Omega^*)$  for any  $\Omega \in S$  ;
- (ii)  $U_1 F(\Omega) + U_1 MG(\Omega) \not\leq_{R_+^s} U_1 F(\Omega^*) + U_1 MG(\Omega^*)$  for any  $\Omega \in S$ .

*Proof.* Let  $Y = \{F(\Omega) + MG(\Omega) : \Omega \in S\}$ . Let  $y = F(\Omega) + MG(\Omega)$  and  $y^* = F(\Omega^*) + MG(\Omega^*)$ . Then  $y^* \notin \mathcal{E}(Y, D)$  if and only if there is a  $y \in Y$  such that  $y^* - y \in D \setminus \{0\}$ . This is so since  $\text{Ker } U_1 = \{0\}$ ,  $y^* - y \in D \setminus \{0\}$  if and only if  $U_1(y^* - y) \geq 0$ . That is,  $U_1 y^* \geq U_1 y$  if and only if  $U_1 y^* \notin \mathcal{E}(U_1 Y, R_+^s)$ .

Now we consider the above lemma for  $D\text{-epi } W$ .

LEMMA 3.3. For  $M \in \mathcal{M}$ , the following are equivalent:

- (i) For some  $(u^*, y^*) \in D\text{-epi } W$  and all  $(u, y) \in D\text{-epi } W$ ,

$$y + Mu \not\leq y^* + Mu^*;$$

- (ii) For some  $\Omega^* \in S$  and all  $\Omega \in S$ ,

$$F(\Omega) + MG(\Omega) \not\leq F(\Omega^*) + MG(\Omega^*).$$

*Proof.* It is similar to that of [7, Lemma 2.5].

Following Nakayama [7], we define a supporting conical variety as follows: For  $M \in \mathcal{M}$ , let  $U_2 = MU_1$  and define a cone  $K$  in  $R^m \times R^p$  by

$$(2) \quad K = \{(u, y) : U_1 y + U_2 u \leq 0, u \in R^m, y \in R^p\}.$$



Let us define the lineality space of  $K$ ,  $K \cup (-K)$ , by  $\ell(K)$ . Since the  $s \times p$  matrix  $U_1$  has the maximal rank  $p$ ,

$$\begin{aligned}\ell(K) &= \{(u, y) : U_1 y + U_2 u = 0\} \\ &= \{(u, y) : y + U_1 u = 0\}.\end{aligned}$$

Since row vectors of  $U_1$  are generators of  $D^\circ$ ,  $MQ \subset D$  implies that every row vector of  $M_2$  is in  $Q^\circ$ .

**DEFINITION 3.4.** We say that  $K$  supports  $D$ -epi  $W$  at  $(u^*, y^*)$  if  $K \cup [cl(D\text{-epi } W) - (u^*, y^*)] \subset \ell(K)$ . We call  $(u^*, y^*)$  the supporting point of  $D$ -epi  $W$ . The set  $\tilde{K} = K + (u^*, y^*)$  is called a conical variety of  $K$ . We also say that  $\tilde{K}$  supports  $D$ -epi  $W$  at  $(u^*, y^*)$ .

The existence of a supporting conical variety of  $D$ -epi  $W$  at a properly efficient point is guaranteed by the following theorem.

**THEOREM 3.5.** Assume Slater's constraint qualification on  $(P)$ . If  $\Omega^*$  is a properly efficient  $D$ -solution to  $(P)$ , then there exists a supporting conical variety  $K$  of  $D$ -epi  $W$  at  $(G(\Omega^*), F(\Omega^*))$ .

*Proof.* Since  $\Omega^*$  is properly efficient, from Theorem 2.1, there exists an  $M^*$  such that

$$(3) \quad F(\Omega) + M^*G(\Omega) \not\leq_D F(\Omega^*) + M^*G(\Omega^*) \text{ for all } \Omega \in \mathcal{S}.$$

If  $(u, y) \in D\text{-epi } W$ , then  $y \in W(u) + D$  so that for some  $\Omega' \in \mathcal{S}(u)$ ,  $F(\Omega') \leq_D y$  and  $G(\Omega') \leq_Q u$ . Let  $K = \{(u, y) : U_1 y + U_1 M^* u \leq 0\}$ , where  $U_1$  is given in (1). Then from (3) and  $F(\Omega') + M^*G(\Omega') \not\leq_D y + M^*u$ , we have

$$y + M^*u \not\leq_D F(\Omega^*) + M^*G(\Omega^*).$$

Therefore, by Lemma 3.2,

$$(4) \quad U_1(y - F(\Omega^*)) + U_2(u - G(\Omega^*)) \not\leq 0.$$

Since  $(u, y) \in D\text{-epi } W$  was arbitrary, (4) holds for all  $(u, y) \in D\text{-epi } W$ . Then, equivalently,  $K$  supports  $D$ -epi  $W$  at  $(G(\Omega^*), F(\Omega^*))$ .



We consider a relationship between a conical variety and supporting hyperplanes. First we define several kinds of half-spaces associated with a hyperplane. Let  $H(\lambda, \mu : r)$  be a hyperplane in  $R^m \times R^p$  with the normal  $(\lambda, \mu)$  such that

$$H(\lambda, \mu : r) = \{(u, y) \in R^m \times R^p : \langle \mu, y \rangle + \langle \lambda, u \rangle - r = 0\}.$$

Associated with the hyperplane  $H(\lambda, \mu : r)$ , half-spaces are defined as follows :

$$H^+(\lambda, \mu : r) = \{(u, y) \in R^m \times R^p : \langle \mu, y \rangle + \langle \lambda, u \rangle - r \geq 0\}$$

$$H_+^o(\lambda, \mu : r) = \{(u, y) \in R^m \times R^p : \langle \mu, y \rangle + \langle \lambda, u \rangle - r > 0\}$$

Similarly,  $H_-$  and  $H_-^o$  are defined by replacing  $\geq$  (respectively,  $=$ ) with  $\leq$  (respectively,  $=$ ).

LEMMA 3.6. [7, Lemma 2.6] *The lineality space of the cone  $K$  defined in (2) with  $U_2 = U_1 M$  is included in the hyperplane  $H(\lambda, \mu : 0)$  if and only if the matrix satisfies  $M^t \mu = \lambda$ .*

Let  $H(\lambda, \mu)$  denote the supporting hyperplane [8, pp.99-100] for  $D$ -epi  $W$  with the inner normal  $(\lambda, \mu)$ , that is,  $H(\lambda, \mu) = H(\lambda, \mu : \tilde{r})$ , where  $\tilde{r} = \sup\{r : H(\lambda, \mu : r) \supset D\text{-epi } W\}$ .

LEMMA 3.7. [7, Lemma 2.7] *For any supporting hyperplane  $H(\lambda, \mu)$  for  $D$ -epi  $W$  at some  $(u, y) \in D\text{-epi } W$ , we have that  $\lambda \in Q^o$  and  $\mu \in D^o$ .*

The next two theorems clarify the relationship between supporting hyperplanes and supporting conical varieties. Note that similar results of Nakayama [7] are not applied directly because the  $D$ -epi  $W$  is not guaranteed to be convex in the programming problem with set functions. Thus we assume more restriction on the perturbed feasible set  $\{\Omega \in \mathcal{S} : G(\Omega) \leq_Q u\}$  in the second theorem.

THEOREM 3.8. *Let  $H(\lambda, \mu)$  be the supporting hyperplane for  $D$ -epi  $W$  with supporting point  $(u^*, y^*)$ . Assume that  $\mu$  is in  $D^o$ . Let  $K$  be a cone defined in (9) for some  $M \in \mathcal{M}$ , and let  $\ell(\tilde{K})$  be the linear*



variety of  $\ell(K)$  passing through  $(u^*, y^*)$ . If  $\ell(\tilde{K})$  is included in  $H(\lambda, \mu)$ , then  $K$  supports  $D$ -epi  $W$  at  $(u^*, y^*)$ .

*Proof.* It is similar to that of [7, Lemma 2.8]. Conversely, given a conical variety of a cone  $K$ , we have a supporting hyperplane of  $D - \text{epi } W$ .

**THEOREM 3.9.** Assume that  $\zeta^o = \zeta$  in the problem (P). If some conical variety  $\tilde{K}$  of a cone  $K$  supports  $D$ -epi  $W$  at  $(u^*, y^*)$ , then there exists a hyperplane  $H(\lambda, \mu : r)$  with  $\mu \neq 0$  supporting  $D$ -epi  $W$  at  $(u^*, y^*)$  such that

$$\ell(\tilde{K}) \subset H(\lambda, \mu : r).$$

*Proof.* Suppose that  $K = \{(u, y) \in R^m \times R^p : U_1 + U_1 M u \leq 0\}$  for some  $M$  and that its conical variety  $K$  supports  $D$ -epi  $W$ , at  $(u^*, y^*)$ . Then  $U_1(y - y^*) + U_1 M(u - u^*) \not\leq 0$  for all  $(u, y) \in D$ -epi  $W$ , or  $y + M u \not\leq_D y^* + M u^*$  for all  $(u, y) \in D$ -epi  $W$ . Note that

$$(5) \quad \begin{aligned} &\{U_1 y + U_1 M u : (u, y) \in D\text{-epi } W\} \\ &\subset \{U_1 y + U_1 M u : (u, y) \in D\text{-epi } \overline{W}\} \end{aligned}$$

where  $\overline{W}(u) = \text{Min}_{D\text{cl}}(F(\mathcal{S}(u)))$  and  $\mathcal{S}(u) = \{\Omega \in \mathcal{S} : G(\Omega) \leq_Q u\}$ . Since  $\zeta^o$  is convex and  $\overline{W}$  is  $D$ -convex, since  $\overline{W}$  is a  $D$ -convex point-to-set map on the convex set  $\zeta^o$ , it follows that the set defined in (5) is a convex set. Using Lemma 3.7 for  $D$ -epi  $\overline{W}$ , we obtain the result. The rest of the proof is similar to that of [7, Lemma 2.8].

## References

1. J.H.Chou, W.S.Hsia, and T.Y.Lee, *Epigraphs of Convex Set Functions*, Journal of Mathematical Analysis and Applications **118** (1986), 247-254.
2. W.S.Hsia, T.Y.Lee, *Lagrangian Functions and Duality Theory in Multiobjective Programming with Set Functions*, Journal of Optimization Theory and Applications **57** (1988), 239-241.
3. W.S.Hsia, T.Y.Lee, and J.Y.Lee, *Lagrange Multiplier Theorem of Multiobjective Programming Problems with Set Functions*, Journal of Optimization Theory and Applications **70** (1991), 137-155.
4. J.Y.Lee, *Weak Duality in Multiobjective Optimization with Set Functions*, Kangweon-Kyungki Mathematical Journal **1** (1993), 65-70.
5. D.G.Luenberger, *Optimization by Vector Space Method*, Wiley, New York, 1969.
6. R.J.T.Morris, *Optimal Constraint Selection of a Measurable Subset*, Journal of Mathematical Analysis and Applications **70** (1979), 546-562.



7. H.Nakayama, *Geometric Consideration of Duality in Vector Optimaization*, Journal of Optimization Theory and Allpications **44** (1984), 625-655.
8. R.T.Rockafellar, *Convex Analysis*, Princeton Univ. Press, Princeton, New Jersey, 1970.
9. Y.Sawaragi, H.Nakayama and T.Tanino, *Theory of Multiobjective Optimization*, Academic Press, Orlando, 1985.

Department of Mathematics Education  
Kangwon National University  
Chunchon, 200-701, Korea



