

## SOME CHARACTERIZATIONS OF KRULL DOMAINS

GYU WHAN CHANG

ABSTRACT. We will find sufficient conditions for a Mori domain to be a Krull domain.

### 1. Introduction

Many of characterizations of Dedekind domains have  $t$ -operation analogues for Krull domains (see [8], [10]). Thus from the well-known characterizations of Dedekind domains, we can deduce new characterizations of Krull domains. In this paper, in spirit of [6, Theorem 37.8 and Theorem 38.1], we find sufficient conditions for a Mori domain to be a Krull domain. In particular, we give examples which show that the conditions of Theorem 6 are the best possible.

Throughout this paper,  $R$  will denote a commutative integral domain with identity and  $K$  its quotient field. Let  $F(R)$  be the set of nonzero fractional ideals of  $R$ . For each  $A \in F(R)$ ,  $A_v = (A^{-1})^{-1}$  and  $A_t = \cup\{J_v : J \text{ is a finitely generated subideal of } A\}$ . If  $A_v = A$  (resp.  $A_t = A$ ) then  $A$  is said to be a divisorial ideal (resp.  $t$ -ideal). We have  $A \subset A_t \subset A_v$ , so that every divisorial ideal is a  $t$ -ideal. If  $A_t = J_t$  for some finitely generated subideal of  $A$ ,  $A_t$  is said to be of finite type.  $R$  is called a Mori domain if each  $t$ -ideal of  $R$  is of finite type, or equivalently, ascending chain condition on  $t$ -ideals holds. By a chain of prime  $t$ -ideals of  $R$  we mean a finite strictly increasing sequence  $P_1 \subsetneq P_2 \subsetneq \dots \subsetneq P_n$ ; the length of the chain is  $n$ . We define the  $t$ -dimension of  $R$ , denoted by  $t\text{-dim}R$ , to be the supremum of the lengths of all chains of prime  $t$ -ideals in  $R$ .

Unexplained terminology is standard, as in [6] or [9].

---

Received March 18, 1998.

1991 Mathematics Subject Classification: 13A15.

Key words and phrases: Krull domain, Mori domain, S-domain,  $t$ -dimension.

## 2. Main results

For a technical reason, we assume that  $R$  is not a field, i.e.,  $R \subsetneq K$ . A maximal  $t$ -ideal of  $R$  is a proper  $t$ -ideal of  $R$  which is maximal among proper  $t$ -ideals of  $R$ . It is easy to see by Zorn's lemma that each maximal  $t$ -ideal is prime and the set of maximal  $t$ -ideals is not empty.

In [1], D. D. Anderson shows that for a proper ideal  $I$  of the ring  $R$  with identity if each prime ideal minimal over  $I$  is finitely generated then the number of prime ideals which are minimal over  $I$  is finite. By the same way as Anderson's proof [1] and the fact that if a prime ideal  $P$  of  $R$  is minimal over a  $t$ -ideal then  $P$  is also a  $t$ -ideal, we have the following useful result.

**LEMMA 1.** *Let  $I$  be a  $t$ -ideal of  $R$ . If each prime ideal  $P$  of  $R$  which is minimal over  $I$  is of finite type, i.e., there is a finitely generated subideal  $A$  of  $P$  such that  $A_t = P$ , then the number of prime ideals minimal over  $I$  is finite.*

**LEMMA 2.** *If  $\{M_\lambda\}$  is the set of maximal  $t$ -ideals of  $R$ , then  $R = \bigcap R_{M_\lambda}$ .*

*Proof.* see [6, Ex. 22, page 52]. □

**DEFINITION 3.** Let  $R$  be an integral domain and  $X^1(R)$  the set of nonzero minimal prime ideals of  $R$ . A domain  $R$  is called a *Krull domain* if

1.  $R = \bigcap_{P \in X^1(R)} R_P$ ,
2.  $R_P$  is a rank one DVR for each  $P \in X^1(R)$ , and
3. for each  $0 \neq a \in R$ , the set of prime ideals of  $X^1(R)$  containing  $a$  is finite.

**THEOREM 4.** (cf. [6, Theorem 38.1]) *Let  $R$  be a Mori domain which is not a field and  $\{M_\alpha\}$  the set of maximal  $t$ -ideals of  $R$ , then the following conditions are equivalent.*

1.  $R$  is a Krull domain.
2. Each  $M_\alpha$  is  $t$ -invertible, i.e.,  $(M_\alpha M_\alpha^{-1})_t = R$ .
3.  $\{(M_\alpha^n)_t\}$  is the set of  $M_\alpha$ -primary ideals and for each  $\alpha$ , there is a prime ideal  $P_\alpha \subsetneq M_\alpha$  such that there are no prime ideals properly between  $P_\alpha$  and  $M_\alpha$ .

*Proof.* (1)  $\implies$  (2) [8, Theorem 3.6].

(2)  $\implies$  (3) [8, Theorem 2.2].

(3)  $\implies$  (1) For a maximal  $t$ -ideal  $M$  of  $R$ ,  $(M^k R_M)_t = ((M^k)_t R_M)_t = (M^k)_t R_M$  [7, Proposition 1.1]. Since  $MR_M$  is a maximal ideal of  $R_M$ ,  $M^k R_M = (MR_M)^k$  is an  $MR_M$ -primary ideal. So  $M^k R_M \cap R$  is  $M$ -primary. So  $M^k R_M \cap R = (M^l)_t$  for some positive integer  $l$ . So  $M^k R_M = (M^k R_M \cap R) R_M = (M^l)_t R_M = (M^l R_M)_t$ . Thus  $M^k R_M = (M^l R_M)_t = ((M^l R_M)_t)_t = (M^k R_M)_t$ . Thus  $\{M^k R_M\}_{k=1}^\infty$  is the set of  $MR_M$ -primary ideals. Let  $P$  be a prime ideal of  $R$  such that  $P \subsetneq M$  and there are no prime ideals properly between  $P$  and  $M$ . Since each  $MR_M/(PR_M)$ -primary ideal is of the form  $A/(PR_M)$  where  $A$  is a  $MR_M$ -primary ideal containing  $PR_M$  and the number of  $MR_M/(PR_M)$ -primary ideals is infinite,  $PR_M \subseteq M^k R_M$  and  $M^k R_M \neq M^{k+1} R_M$  for each positive integer  $k$ . Since  $R_M$  is a Mori domain,  $PR_M = 0$  [8, Theorem 2.1] and hence  $P = 0$ . Thus  $R_M$  is a rank one DVR. By Lemmas 1 and 2,  $R$  is a Krull domain.  $\square$

A domain  $R$  is said to be a Prüfer  $v$ -multiplication domain (PVMD) if each finitely generated ideal  $I$  of  $R$  is  $t$ -invertible, i.e.,  $(II^{-1})_t = R$ , or equivalently,  $R_P$  is a valuation domain, for each maximal  $t$ -ideal  $P$ . Recall from [9, page 26] that a domain  $R$  is called an  $S$ -domain if for every height one prime ideal  $P$ , the expansion  $P[X]$  of  $P$  to the polynomial ring  $R[X]$  also has height one.

LEMMA 5. *Let  $R$  be a domain of  $t\text{-dim} R \leq 1$ , then  $R$  is an integrally closed  $S$ -domain if and only if  $R$  is a PVMD.*

*Proof.* ( $\implies$ ) If  $P$  is a maximal  $t$ -ideal,  $R_P$  is an one dimensional integrally closed domain. Since  $R$  is an  $S$ -domain,  $ht(P[X]) = 1$ . So  $dim R_P[X] = dim(R_P[X]_{PR_P[X]}) + 1 = dim(R[X]_{P[X]}) + 1 = ht(P[X]) + 1 = 2$ . Since  $R_P$  is integrally closed,  $R_P$  is a valuation domain [6, Proposition 30.14].

( $\impliedby$ ) If  $P$  is a prime ideal of  $ht(P) = 1$ ,  $R_P$  is a valuation domain. So  $dim(R_P[X]) = 2$  and hence  $ht(P[X]) = ht(PR_P[X]) = 1$ .  $\square$

THEOREM 6. (cf. [6, Theorem 37.8]) *A domain  $R$  is a Krull domain if (and only if)  $R$  is an integrally closed Mori domain of  $t\text{-dim} R = 1$  and  $R$  is an  $S$ -domain.*

*Proof.* By Lemma 5,  $R$  is a PVMD. Since  $R$  is a Mori domain,  $R_P$  is a rank one DVR for each maximal  $t$ -ideal  $P$ . By Lemmas 1 and 2,  $R$  is a Krull domain.  $\square$

In Theorem 6, the hypothesis that  $R$  is an  $S$ -domain is necessary. To see this, we give an example of an integrally closed Mori domain  $R$  of  $t\text{-dim}R = 1$ , which is not a Krull domain.

**EXAMPLE 7.** Let  $C$  (resp.  $Q$ ) be the field of complex (resp. rational) numbers and  $\overline{Q}$  the algebraic closure of  $Q$  in  $C$ . Then the subring  $D = \overline{Q} + XC[[X]]$  of the power series ring  $C[[X]]$  is an integrally closed Mori domain of  $\dim D = 1$  [4, Theorem 3.2]. But  $D$  is not a valuation domain [5, Theorem 2.1(h)]. Thus  $D$  is not a Krull domain.

**EXAMPLE 8.** Let  $\mathfrak{R}$  be the field of real numbers and  $R = \mathfrak{R}[[x, y]] = \mathfrak{R} + M$ , where  $M = (x, y)$ , the power series ring over  $\mathfrak{R}$ . Let  $\overline{Q}$  be the algebraic closure of the field  $Q$  of rational numbers in  $\mathfrak{R}$ . Let  $D = \overline{Q} + M$ , then

1.  $D$  is integrally closed,
2.  $D$  is a Mori domain of  $t\text{-dim}D = 2$  and  $D$  satisfies Krull's principal ideal theorem and
3.  $D$  is an  $S$ -domain.

*Proof.* 1. Since  $\overline{Q}$  is the integral closure of  $Q$  in  $\mathfrak{R}$ ,  $D$  is integrally closed [5, Theorem 2.1.(b)].

2. By [9, Theorem 71 and Theorem 72],  $R$  is a Noetherian UFD of  $\dim R = 2$  and  $M$  is the unique maximal ideal of  $R$ . By [2, Proposition 3.8],  $\text{Spec}(R) = \text{Spec}(D)$ . So  $D$  is a Mori domain [4, Theorem 3.2] and  $D$  satisfies Krull's principal ideal theorem [3, Corollary 3.2]. By [2, Proposition 3.23],  $M$  is a  $t$ -ideal of  $D$ . Since  $ht(M) = 2$ ,  $t\text{-dim}R = 2$ .

3. If  $P$  is a prime ideal of  $D$  of  $ht(P) = 1$ ,  $R \subseteq D_P$ . For if  $m \in M - P$ ,  $r = (rm)\frac{1}{m} \in D_P$  for each  $r \in R$ . Since  $\text{Spec}(R) = \text{Spec}(D)$ ,  $R_P \subseteq (D_P)_{(R-P)} = D_P$ . Since  $R_P$  is a rank one DVR,  $R_P = D_P$ . Thus the polynomial ring  $D_P[X] = R_P[X]$  is of dimension 2, and  $ht(P[X]) = ht(PD_P[X]) = 1$ . So  $D$  is an  $S$ -domain.  $\square$

Example 7 and Example 8 show that the hypothesis in Theorem 6 for an integrally closed Mori domain to be a Krull domain cannot be weakened.

REMARK 9. In Theorem 6, the condition that  $R$  is a Mori domain can be replaced by the assumption that each prime  $t$ -ideal is of finite type. For if  $P$  is a prime  $t$ -ideal, there is a finitely generated subideal  $I$  of  $P$  such that  $P = I_t$ . So  $PR_P = I_tR_P = (I_tR_P)_t = (IR_P)_t$ . Since  $R_P$  is a valuation domain,  $PR_P$  is principal and so  $R_P$  is a local PID.

### References

- [1] D. D. Anderson, *A note on minimal prime ideals*, Proc. Amer. Math. Soc. 122(1), (1994), 13-14.
- [2] D. F. Anderson and D. E. Dobbs, *Pairs of rings with the same prime ideals*, Canad. J. Math. 32(1980), 362 - 384.
- [3] V. Barucci, D. F. Anderson and D. E. Dobbs, *Coherent Mori domain and the principal ideal theorem*, Comm. Algebra 15(6) (1987) 1119 - 1156.
- [4] V. Barucci, *On a class of Mori domains*, Comm. Algebra 11(17) (1983) 1989 -2001.
- [5] E. Bastida and R. Gilmer, *Overrings and divisorial ideals of the form  $D + M$* , Michigan. Math. J. 20(1973) 79 - 95.
- [6] R. Gilmer, *Multiplicative ideal Theory*, Marcel Dekker, New York, 1972.
- [7] E. G. Houston, T. G. Lucas and T. M. Viswanathan, *Primary decomposition of divisorial ideals in Mori domains*, J. Algebra 117 (1988) 327 - 342.
- [8] B. G. Kang, *A converse of the well known fact about Krull domains*, J. Algebra 124 (1989) 284 - 299.
- [9] I. Kaplansky, *Commutative Rings, The Revision*, Univ. of Chicago, Chicago, 1974.
- [10] S. Malik, J. L. Matt and M. Zafrullah, *On  $t$ -invertibility*, Comm. Algebra 16(1) (1988) 149 - 170.

Department of Mathematics  
College of Natural Science, Kangwon National University  
Chuncheon, 200-701, Korea