

CONVERGENCE OF EXPONENTIALLY BOUNDED C -SEMIGROUPS

YOUNG S. LEE

ABSTRACT. In this paper, we establish the conditions that a mild C -existence family yields a solution to the abstract Cauchy problem. And we show the relation between mild C -existence family and C -regularized semigroup if the family of linear operators is exponentially bounded and C is a bounded injective linear operator.

1. Introduction

Let X be a Banach space and let A be a linear operator from $D(A) \subset X$ into X . The abstract Cauchy problem for A with initial data $x \in X$ is to find a solution $u(t)$ to the following initial value problem

$$\frac{du}{dt} = Au, \quad t > 0, \quad u(0) = x. \quad (\text{ACP})$$

It is well known ([5]) that when A is closed, generating a C_0 semigroup $\{T(t) : t \geq 0\}$ guarantees the abstract Cauchy problem to have a unique mild solution for all initial data $x \in X$, and the solution is given by $u(t) = T(t)x$.

If the abstract Cauchy problem does not have a mild solution for all $x \in X$, we may look for initial data in X that produce mild solutions. In [4], a family of linear operators was introduced that produces a solution of the abstract Cauchy problem for all initial data in the range of a bounded linear operator C . It is known [4] that for a bounded linear operator C mild C -existence family yields a mild solution for all initial data in the range of C . C -existence family is a generalization of the classical C_0 semigroup such as integrated semigroups and regularized semigroups (see [1, 2, 3, 4, 6]).

Received July 12, 2001.

2000 Mathematics Subject Classification: Primary 47D05, Secondary 47D10.

Key words and phrases: Existence Family, C -semigroups, mild solution.

In this paper, we establish the conditions that mild C -existence family for A produces a solution of (ACP). It is known that a mild C -existence family for A is a C -regularized semigroup generated by an extension of A if the family of operators and A commute. We will show the relation between C -regularized semigroup and mild C -existence family if the family of operators is exponentially bounded. If $CA \subset AC$, mild C -existence family is a C -regularized semigroup generated by an extension of A .

Throughout this paper, we denote by $D(A)$ the domain of the operator A on X and $R(A)$ the range of A . An operator T in X is said to commute with A if $AT \subset TA$. It means that if $x \in D(T)$, $Ax \in D(T)$ and $TAx = ATx$. By a solution $u(t)$ of (ACP), we mean a continuously differentiable function $u : [0, \infty) \rightarrow X$ such that $u(t) \in D(A)$ for all $t \geq 0$ and satisfies (ACP). A mild solution of (ACP) is a continuous function $u : [0, \infty) \rightarrow X$ such that $v(t) = \int_0^t u(s)ds \in D(A)$ and

$$\frac{d}{dt}v(t) = Av(t) + x, \quad t \geq 0.$$

section2. Mild C -existence Families

Let A be as in (ACP).

DEFINITION 2.1. The strongly continuous family $\{S(t) : t \geq 0\}$ of bounded linear operators on X is called a mild C -existence family for A if for all $x \in X$, $t > 0$, $\int_0^t S(s)xds \in D(A)$ and

$$A \left(\int_0^t S(s)xds \right) = S(t)x - Cx.$$

REMARK. If $\{S(t) : t \geq 0\}$ is a mild C -existence family for A , then $u(t) = S(t)x$ is a mild solution of (ACP) with $u(0) = Cx$.

THEOREM 2.2. Let $\{S(t) : t \geq 0\}$ be a mild C -existence family for A . Suppose that $CA \subset AC$. Then there exists a solution $u(t)$ of (ACP) with $u(0) = x \in C(D(A))$.

Proof. Let $x \in C(D(A))$. Then there exists $y \in D(A)$ such that $x = Cy$. So $Ax = ACy = CAy$. Define

$$u(t) = x + \int_0^t S(s)Ayds.$$

Then $du(t)/dt = S(t)Ay$ and $Au(t) = Ax + A(\int_0^t S(s)Ayds) = Ax + S(t)Ay - CAy = S(t)Ay$, since $\{S(t) : t \geq 0\}$ is a mild C -existence family for A . So $u(t)$ is a solution of (ACP) with $u(0) = x$. \square

THEOREM 2.3. *Let $\{S(t) : t \geq 0\}$ be a mild C -existence family for A . If $x \in D(A)$ and $Ax \in R(C)$, then there exists a solution $u(t)$ for (ACP) with initial data x .*

Proof. Let $y \in X$ such that $Ax = Cy$. Define

$$u(t) = x + \int_0^t S(s)yds.$$

Then $du(t)/dt = S(t)y$ and $Au(t) = Ax + A \int_0^t S(s)yds = Ax + S(t)y - Cy = S(t)y$, since $\{S(t) : t \geq 0\}$ is a mild C -existence family for A . Thus $u(t)$ is a solution of (ACP) with $u(0) = x$. \square

Next, we characterize an exponentially bounded mild C -existence family in terms of Laplace transforms.

THEOREM 2.4. *Let $\{S(t) : t \geq 0\}$ be a strongly continuous family of bounded linear operators on X such that $\|S(t)\| \leq Me^{\omega t}$, $t \geq 0$, for some M , $\omega > 0$. Suppose that A is closed and $\lambda - A$ is injective for $\lambda > \omega$. Then the following are equivalent.*

- (1) $\{S(t) : t \geq 0\}$ is a mild C -existence family for A .
- (2) $R(C) \subset R(\lambda - A)$ for $\lambda > \omega$ and

$$(\lambda - A)^{-1}Cx = \int_0^\infty e^{-\lambda t} S(t)xdt, \quad x \in X, \lambda > \omega.$$

Proof. Suppose that $\{S(t) : t \geq 0\}$ is a mild C -existence family for A . Let $x \in X$. By integration by parts, we have

$$\int_0^\infty e^{-\lambda t} S(t)xdt = \lambda \int_0^\infty e^{-\lambda t} \left(\int_0^t S(s)xds \right) dt.$$

Since A is closed,

$$\begin{aligned}
 A \int_0^\infty e^{-\lambda t} S(t)x dt &= \lambda A \int_0^\infty e^{-\lambda t} \left(\int_0^t S(s)x ds \right) dt \\
 &= \lambda \int_0^\infty e^{-\lambda t} A \left(\int_0^t S(s)x ds \right) dt \\
 &= \lambda \int_0^\infty e^{-\lambda t} (S(t)x - Cx) dt \\
 &= \lambda \int_0^\infty e^{-\lambda t} S(t)x dt - Cx.
 \end{aligned}$$

So $Cx = (\lambda - A) \int_0^\infty e^{-\lambda t} S(t)x dt$ for all $x \in X$, and the result follows.

Suppose that (2) is satisfied. Let $x \in X$. By Post-Widder inversion theorem of Laplace transform and the closedness of A , we have

$$\begin{aligned}
 \lambda^{-1} A (\lambda - A)^{-1} Cx &= \lambda^{-1} A \int_0^\infty e^{-\lambda t} S(t)x dt \\
 &= \int_0^\infty e^{-\lambda t} A \left(\int_0^t S(s)x ds \right) dt.
 \end{aligned}$$

Since $Cx = (\lambda - A)(\lambda - A)^{-1} Cx$,

$$\begin{aligned}
 \lambda^{-1} A (\lambda - A)^{-1} Cx &= (\lambda - A)^{-1} Cx - \lambda^{-1} Cx \\
 &= \int_0^\infty e^{-\lambda t} (S(t)x - Cx) dt.
 \end{aligned}$$

By the uniqueness of the Laplace transform, we have

$$A \left(\int_0^t S(s)x ds \right) = S(t)x - Cx. \quad \square$$

The following shows the relation between mild C -existence family and regularized semigroup.

THEOREM 2.5. *Let $\{S(t) : t \geq 0\}$ be an exponentially bounded mild C -existence family for A such that $\|S(t)\| \leq M e^{\omega t}$ for $t \geq 0$, some M and $\omega > 0$, and let C be a bounded injective linear operator. Suppose that A is closed and has no eigenvalue in (ω, ∞) with $CA \subset AC$. Then $\{S(t) : t \geq 0\}$ is a C -regularized semigroup generated by an extension of A .*

Proof. Let $x \in X$. Since $\{S(t) : t \geq 0\}$ is a mild C -existence family and $CA \subset AC$,

$$CS(t)x - C^2x = CA \left(\int_0^t S(s)x ds \right) = A \left(\int_0^t CS(s)x ds \right).$$

So $CS(t)x$ is a mild solution of (ACP) with $u(0) = C^2x$. Clearly, $S(t)Cx$ is a mild solution of (ACP) with $u(0) = C^2x$. By the uniqueness of mild solution (Proposition in [4]), $CS(t) = S(t)C$.

By Theorem 2.4, for $x \in X$,

$$\begin{aligned} C(\lambda - A)^{-1}Cx &= C \int_0^\infty e^{-\lambda t} S(t)x dt \\ &= \int_0^\infty e^{-\lambda t} S(t)Cx dt = (\lambda - A)^{-1}C^2x. \end{aligned}$$

So we have

$$\begin{aligned} &(\lambda - A)^{-1}C(\mu - A)^{-1}Cx \\ &= \frac{1}{\lambda}A(\lambda - A)^{-1}C(\mu - A)^{-1}Cx + \frac{1}{\lambda}C(\mu - A)^{-1}Cx \\ &= \frac{1}{\lambda}(\lambda - A)^{-1}CA(\mu - A)^{-1}Cx + \frac{1}{\lambda}(\mu - A)^{-1}C^2x \\ &= \frac{1}{\lambda}(\lambda - A^{-1})C(\mu(\mu - A)^{-1}Cx - Cx) + \frac{1}{\lambda}(\mu - A)^{-1}C^2x \\ &= \frac{\mu}{\lambda}(\lambda - A)^{-1}C(\mu - A)^{-1}Cx - \frac{1}{\lambda}(\lambda - A)^{-1}C^2x + \frac{1}{\lambda}(\mu - A)^{-1}C^2x \end{aligned}$$

Thus

$$(\lambda - \mu)(\lambda - A)^{-1}C(\mu - A)^{-1}Cx = (\mu - A)^{-1}C^2x - (\lambda - A)^{-1}C^2x.$$

By Theorem 2.4 and integration by parts,

$$\begin{aligned}
& \frac{1}{\lambda - \mu} ((\mu - A)^{-1}C^2x - (\lambda - A)^{-1}C^2x) \\
&= \int_0^\infty e^{-(\lambda-\mu)t} (\mu - A)^{-1}C^2x dt - \frac{1}{\lambda - \mu} \int_0^\infty e^{-\lambda t} S(t)Cx dt \\
&= \int_0^\infty e^{-(\lambda-\mu)t} (\mu - A)^{-1}C^2x dt - \int_0^\infty \frac{1}{\lambda - \mu} e^{-(\lambda-\mu)t} e^{-\mu t} S(t)Cx dt \\
&= \int_0^\infty e^{-(\lambda-\mu)t} \left(\int_0^\infty e^{-\mu s} S(s)Cx ds \right) dt \\
&\quad - \int_0^\infty e^{-(\lambda-\mu)t} \left(\int_0^t e^{-\mu s} S(s)Cx ds \right) dt \\
&= \int_0^\infty e^{-(\lambda-\mu)t} \left(\int_t^\infty e^{-\mu s} S(s)Cx ds \right) dt \\
&= \int_0^\infty e^{-\lambda t} \left(\int_t^\infty e^{-\mu(s-t)} S(s)Cx ds \right) dt \\
&= \int_0^\infty e^{-\lambda t} \left(\int_0^\infty e^{-\mu w} S(t+w)Cx dw \right) dt
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
(\lambda - A)^{-1}C(\mu - A)^{-1}Cx &= \int_0^\infty e^{-\lambda t} S(t)(\mu - A)^{-1}Cx dt \\
&= \int_0^\infty e^{-\lambda t} S(t) \left(\int_0^\infty e^{-\mu s} S(s)Cx ds \right) dt \\
&= \int_0^\infty e^{-\lambda t} \left(\int_0^\infty e^{-\mu s} S(t)S(s)Cx ds \right) dt.
\end{aligned}$$

By the uniqueness of the Laplace transform, we have

$$S(t+s)C = S(t)S(s).$$

Therefore $\{S(t) : t \geq 0\}$ is a C -regularized semigroup generated by an extension of A . \square

References

- [1] W. Arendt, *Vector valued Laplace transform and Cauchy problems*, Israel J. Math. **59** (1987), 327 - 352.
- [2] E.B. Davies and M.M.H. Pang, *The Cauchy problem and a generalization of the Hille-Yosida theorem*, Proc. London Math. Soc. **55** (1987), 181 - 248.
- [3] R. deLaubenfels, *C -semigroups and the Cauchy problem*, J. Funct. Anal. **111** (1993), 44 - 61.
- [4] R. deLaubenfels, *Existence families, functional calculi and evolution equations*, Lecture Notes in Math., vol. 1570, Springer Verlag, 1994.
- [5] J. A. Goldstein, *Semigroups of operators and applications*, Oxford, New York, 1985.
- [6] M. Li and F. Huang, *Characterization of contraction C -semigroups*, Proc. Amer. Math. Soc. **126** (1998), 1063 - 1069.

Department of Mathematics
Seoul Women's University
Seoul, 139 - 774, Korea
E-mail: younglee@swift2.swu.ac.kr