

CONFORMAL CHANGE OF THE VECTOR S_ω IN 5-DIMENSIONAL g -UFT

CHUNG HYUN CHO

ABSTRACT. We investigate change of the vector S_ω induced by the conformal change in 5-dimensional g -unified field theory. These topics will be studied for the second class in 5-dimensional case

1. Introduction

The conformal change in a generalized 4-dimensional Riemannian space connected by an Einstein's connection was primarily studied by HLAVATÝ ([8], 1957). CHUNG ([6], 1968) also investigated the same topic in 4-dimensional $*g$ -unified field theory.

The Einstein's connection induced by the conformal change for all classes in 3-dimensional case, for the second and third classes in 5-dimensional case, and for the first class in 5-dimensional $*g$ -UFT, and for the second class in 5-dimensional g -UFT were investigated by CHO ([1], 1992, [2], 1994, [3], 1996, [4], 1998).

In the present paper, we investigate change of the vector S_ω induced by the conformal change in 5-dimensional g -unified field theory. These topics will be studied for the second class in 5-dimensional case.

2. Preliminaries

This chapter is a brief collection of basic concepts, notations, theorems, and results needed in our further considerations. They may be referred to CHUNG ([5], 1988; [3], 1988), CHO ([1], 1992; [2], 1994; [3], 1996; [4], 1998).

Received November 19, 2002.

2000 Mathematics Subject Classification: 53A45, 83E99.

Key words and phrases: Einstein connection, conformal change.

2.1. n -dimensional g -unified field theory. The n -dimensional g -unified field theory (n - g -UFT hereafter) was originally suggested by HLAVATÝ([8],1957) and systematically introduced by CHUNG([7],1963).

Let X_n ¹ be an n -dimensional generalized Riemannian manifold, referred to a real coordinate system x^ν obeying coordinate transformations $x^\nu \rightarrow x^{\nu'}$, for which

$$(2.1) \quad \text{Det}\left(\left(\frac{\partial x}{\partial x'}\right)\right) \neq 0.$$

In the usual Einstein's n -dimensional unified field theory, the manifold X_n is endowed with a general real nonsymmetric tensor $g_{\lambda\mu}$ which may be split into its symmetric part $h_{\lambda\mu}$ and skew-symmetric part $k_{\lambda\mu}$ ²:

$$(2.2) \quad g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu}$$

where

$$(2.3) \quad \text{Det}((g_{\lambda\mu})) \neq 0 \quad \text{Det}((h_{\lambda\mu})) \neq 0.$$

Therefore we may define a unique tensor $h^{\lambda\nu} = h^{\nu\lambda}$ by

$$(2.4) \quad h_{\lambda\mu} h^{\lambda\nu} = \delta_\mu^\nu.$$

In our n - g -UFT, the tensors $h_{\lambda\mu}$ and $h^{\lambda\nu}$ will serve for raising and/or lowering indices of the tensors in X_n in the usual manner.

The manifold X_n is connected by a general real connection $\Gamma_{\omega\mu}^\nu$ with the following transformation rule :

$$(2.5) \quad \Gamma_{\omega'\mu'}^{\nu'} = \frac{\partial x^{\nu'}}{\partial x^\alpha} \left(\frac{\partial x^\beta}{\partial x^{\omega'}} \cdot \frac{\partial x^\gamma}{\partial x^{\mu'}} \Gamma_{\beta\gamma}^\alpha + \frac{\partial^2 x^\alpha}{\partial x^{\omega'} \partial x^{\mu'}} \right)$$

and satisfies the system of Einstein's equations

$$(2.6) \quad D_\omega g_{\lambda\mu} = 2S_{\omega\mu}{}^\alpha g_{\lambda\alpha}$$

where D_ω denotes the covariant derivative with respect to $\Gamma_{\lambda\mu}^\nu$ and

$$(2.7) \quad S_{\lambda\mu}{}^\nu = \Gamma_{[\lambda\mu]}^\nu$$

is the *torsion tensor* of $\Gamma_{\lambda\mu}^\nu$. The connection $\Gamma_{\lambda\mu}^\nu$ satisfying (2.6) is called the *Einstein's connection*.

¹Throughout the present paper, we assumed that $n \geq 2$.

²Throughout this paper, Greek indices are used for holonomic components of tensors. In X_n all indices take the values $1, \dots, n$ and follow the summation convention.

In our further considerations, the following scalars, tensors, abbreviations, and notations for $p = 0, 1, 2, \dots$ are frequently used :

$$\begin{aligned} \mathfrak{g} &= \text{Det}((g_{\lambda\mu})) \neq 0, & \mathfrak{h} &= \text{Det}((h_{\lambda\mu})) \neq 0, \\ \mathfrak{t} &= \text{Det}((k_{\lambda\mu})), \end{aligned} \quad (2.8a)$$

$$g = \frac{\mathfrak{g}}{\mathfrak{h}}, \quad k = \frac{\mathfrak{t}}{\mathfrak{h}}, \quad (2.8b)$$

$$K_p = k_{[\alpha_1}^{\alpha_1} \cdots k_{\alpha_p]}^{\alpha_p}, \quad (p = 0, 1, 2, \dots) \quad (2.8c)$$

$${}^{(0)}k_\lambda^\nu = \delta_\lambda^\nu, \quad {}^{(1)}k_\lambda^\nu = k_\lambda^\nu, \quad {}^{(p)}k_\lambda^\alpha = {}^{(p-1)}k_\lambda^\alpha k_\alpha^\nu, \quad (2.8d)$$

$$K_{\omega\mu\nu} = \nabla_\nu k_{\omega\mu} + \nabla_\omega k_{\nu\mu} + \nabla_\mu k_{\omega\nu}, \quad (2.8e)$$

$$\sigma = \begin{cases} 1 & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases} \quad (2.8f)$$

where ∇_ω is the symbolic vector of the covariant derivative with respect to the Christoffel symbols $\{\lambda_\mu\}$ defined by $h_{\lambda\mu}$. The scalars and vectors introduced in (2.8) satisfy

$$K_0 = 1; K_n = k \quad \text{if } n \text{ is even}; \quad K_p = 0 \quad \text{if } p \text{ is odd}, \quad (2.9a)$$

$$g = 1 + K_2 + \cdots + K_{n-\sigma}, \quad (2.9b)$$

$${}^{(p)}k_{\lambda\mu} = (-1)^{p(p)} k_{\mu\lambda}, \quad {}^{(p)}k^{\lambda\mu} = (-1)^{p(p)} k^{\nu\lambda}. \quad (2.9c)$$

Furthermore, we also use the following useful abbreviations, denoting an arbitrary tensor $T_{\omega\mu\nu}$, skew-symmetric in the first two indices, by T:

$${}^{pqr}T = {}^{pqr}T_{\omega\mu\nu} = T_{\alpha\beta\gamma} {}^{(p)}k_\omega^{\alpha(q)} k_\mu^{\beta(r)} k_\nu^\gamma, \quad (2.10a)$$

$$T = T_{\omega\mu\nu} = {}^{000}T, \quad (2.10b)$$

$$2 {}^{pqr}T_{\omega[\lambda\mu]} = {}^{pqr}T_{\omega\lambda\mu} - {}^{pqr}T_{\omega\mu\lambda}, \quad (2.10c)$$

$$2 {}^{(pq)r}T_{\omega\lambda\mu} = {}^{pqr}T_{\omega\lambda\mu} + {}^{qpr}T_{\omega\lambda\mu}. \quad (2.10d)$$

We then have

$${}^{pqr}T_{\omega\lambda\mu} = -{}^{qpr}T_{\lambda\omega\mu}. \quad (2.11)$$

If the system (2.6) admits $\Gamma_{\lambda\mu}^\nu$, using the above abbreviations it was shown that the connection is of the form

$$\Gamma_{\omega\mu}^\nu = \{\omega\mu\}^\nu + S_{\omega\mu}^\nu + U_{\omega\mu}^\nu \quad (2.12)$$

where

$$U_{\nu\omega\mu} = 2 S_{\nu(\omega\mu)}. \quad (2.13)$$

The above two relations show that our problem of determining $\Gamma_{\omega\mu}^\nu$ in terms of $g_{\lambda\mu}$ is reduced to that of studying the tensor $S_{\omega\mu}^\nu$. On the other hand, it has also been shown that the tensor $S_{\omega\mu}^\nu$ satisfies

$$S = B - 3 \overset{(110)}{S} \quad (2.14)$$

where

$$2B_{\omega\mu\nu} = K_{\omega\mu\nu} + 3K_{\alpha[\mu\beta}k_{\omega]}^\alpha k_\nu^\beta. \quad (2.15)$$

DEFINITION 2.1. The vector S_ω defined by

$$S_\omega = S_{\omega\alpha}^\alpha. \quad (2.16)$$

2.2. Some results for the second class in 5-g-UFT. In this section, we introduce some results of 5-g-UFT without proof, which are needed in our subsequent considerations.

They may be referred to CHO([1],1992).

DEFINITION 2.2. In 5-g-UFT, the tensor $g_{\lambda\mu}(k_{\lambda\mu})$ is said to be the second class, if $K_2 \neq 0$, $K_4 = 0$.

THEOREM 2.3 (Main Recurrence Relations). *For the second class in 5-UFT, the following recurrence relation hold*

$$\overset{(p+3)}{k_\lambda}{}^\nu = -K_2^{\overset{(p+1)}{k_\lambda}{}^\nu}, \quad (p = 0, 1, 2, \dots). \quad (2.17)$$

THEOREM 2.4 (For The Second Class In 5-g-UFT). *A necessary and sufficient condition for the existence and uniqueness of the solution of (2.5) is*

$$1 - (K_2)^2 \neq 0. \quad (2.18)$$

If the condition (2.18) is satisfied, the unique solution of (2.14) is given by

$$(1 - K_2^2)(S - B) = -2 \overset{(10)1}{B} + (K_2 - 1) \overset{110}{B} + 2 \overset{(20)2}{B} + 2 \overset{112}{B}. \quad (2.19)$$

3. Conformal change of the 5-dimensional vector S_ω for the second class

In this final chapter we investigate the change $S_\omega \rightarrow \bar{S}_\omega$ of the vector induced by the conformal change of the tensor $g_{\lambda\mu}$, using the recurrence relations and theorems introduced in the preceding chapter.

We say that X_n and \bar{X}_n are conformal if and only if

$$\bar{g}_{\lambda\mu}(x) = e^\Omega g_{\lambda\mu}(x) \quad (3.1)$$

where $\Omega = \Omega(x)$ is an at least twice differentiable function. This conformal change enforces a change of the vector S_ω . An explicit representation of the change of 5-dimensional vector S_ω for the second class will be exhibited in this chapter.

AGREEMENT 3.1. Throughout this section, we agree that, if T is a function of $g_{\lambda\mu}$, then we denote \bar{T} the same function of $\bar{g}_{\lambda\mu}$. In particular, if T is a tensor, so is \bar{T} . Furthermore, the indices of $T(\bar{T})$ will be raised and/or lowered by means of $h^{\lambda\nu}(\bar{h}^{\lambda\nu})$ and/or $h_{\lambda\nu}(\bar{h}_{\lambda\nu})$.

The results in the following theorems are needed in our further considerations. They may be referred to CHO([1],1992, [2],1994, [3],1996).

THEOREM 3.2. *In n - g -UFT, the conformal change (3.1) induces the following changes:*

$${}^{(p)}\bar{k}_{\lambda\mu} = e^{\Omega(p)} k_{\lambda\mu}, \quad {}^{(p)}\bar{k}_\lambda = {}^{(p)}k_\lambda^\nu, \quad {}^{(p)}\bar{k}^{\lambda\mu} = e^{-\Omega(p)} k^{\lambda\mu}, \quad (3.2a)$$

$$\bar{g} = g, \quad \bar{K}_p = K_p, \quad (p = 1, 2, \dots). \quad (3.2b)$$

THEOREM 3.3 (For all classes in 5- g -UFT). *The change of the tensor $B_{\omega\mu\nu}$ induced by the conformal change (3.1) may be given by*

$$\begin{aligned} \bar{B}_{\omega\mu\nu} &= e^\Omega (B_{\omega\mu\nu} + k_{\nu[\omega} \Omega_{\mu]} - k_{\omega\mu} \Omega_\nu \\ &\quad - h_{\nu[\omega} k_{\mu]}^\delta \Omega_\delta + 2^{(2)} k_{\nu[\omega} k_{\mu]}^\delta \Omega_\delta + k_{\omega\mu} {}^{(2)} k_{\nu}^\delta \Omega_\delta). \end{aligned} \quad (3.3)$$

Now, we are ready to derive representations of the changes $S_\omega \rightarrow \bar{S}_\omega$ in 5- g -UFT for the second class induced by the conformal change (3.1).

THEOREM 3.4. *The conformal change (3.1) induces the following change :*

$$\begin{aligned} \overline{{}^{(10)1} B}_{\omega\mu\nu} &= e^\Omega [{}^{(10)1} B_{\omega\mu\nu} + (-2^{(4)} k_{\nu[\omega} k_{\mu]}^\delta \\ &\quad + 2^{(2)} k_{\nu[\omega} k_{\mu]}^\delta - k_{\nu[\omega} {}^{(2)} k_{\mu]}^\delta) \Omega_\delta - {}^{(3)} k_{\nu[\omega} \Omega_{\mu]}]. \end{aligned} \quad (3.4)$$

THEOREM 3.5. *The conformal change (3.1) induces the following change:*

$$\begin{aligned} \overline{{}^{ppq} B}_{\omega\mu\nu} &= e^\Omega [{}^{ppq} B_{\omega\mu\nu} + (-1)^p \{ 2^{(p+q+2)} k_{\nu[\omega} {}^{(p+1)} k_{\mu]}^\delta \\ &\quad + {}^{(2p+1)} k_{\omega\mu} {}^{(2+q)} k_{\nu}^\delta - (2p+1) k_{\omega\mu} {}^{(q)} k_{\nu}^\delta \\ &\quad + {}^{(p+q+1)} k_{\nu[\omega} {}^{(p)} k_{\mu]}^\delta - (p+q) k_{\nu[\omega} {}^{(p+1)} k_{\mu]}^\delta \} \Omega_\delta]. \end{aligned} \quad (3.5)$$

$$\left(\begin{array}{l} p = 0, 1, 2, 3, 4, \dots \\ q = 0, 1, 2, 3, 4, \dots \end{array} \right)$$

THEOREM 3.6. *The change $S_\omega \rightarrow \bar{S}_\omega$ induced by conformal change (3.1) may be represented by*

$$\begin{aligned} \bar{S}_\omega &= S_\omega + \frac{1}{2C} [(-7K_2^3 + 11K_2^2 - 8K_2 + 1)k_\omega^\delta \Omega_\delta \\ &\quad - (3 - K_2 + K_2^2)^{(2)} k_\alpha^\alpha k_\omega^\delta \Omega_\delta] \end{aligned} \quad (3.6)$$

where $C = K_2^2 - 1$.

Proof. In virtue of (2.19) and Agreement (3.1), we have

$$(1 - \bar{K}_2^2)(\bar{S} - \bar{B}) = -2 \frac{\overline{(10)1}}{B} + (\bar{K}_2 - 1) \frac{\overline{110}}{B} + 2 \frac{\overline{(20)2}}{B} + 2 \frac{\overline{112}}{B}. \quad (3.7)$$

The relation(3.6) follows by substituting (3.2), (3.3), (3.4), (3.5), (2.17), Definition (2.1), Definition (2.2) into (3.7). \square

REMARK 3.7. The results (3.6) can also be obtained by using Theorem 3.6 of [4] and (2.16).

References

- [1] C.H. Cho, *Conformal change of the connection in 3- and 5-dimensional $*g^{\lambda\nu}$ -unified Field Theory*, BIBS, Inha Univ. **13** (1992).
- [2] C.H. Cho, *Conformal change of the connection for the first class in 5-dimensional $*g^{\lambda\nu}$ -unified Field Theory*, Kangweon-Kyungki Math. Jour. **2** (1994), 19-33.
- [3] C.H. Cho, *Conformal change of the tensor $S_{\lambda\mu}{}^\nu$ for the second category in 6-dimensional g -UFT*, Kangweon-Kyungki Math. Jour. **4** (1996), 163-171.
- [4] C.H. Cho, *Conformal change of the torsion tensor in 5-dimensional g -unified field theory*, Kangweon-Kyungki Math. Jour. **6**, No.2 (1998), 213-220.
- [5] K.T. Chung, *Three- and Five-dimensional considerations of the geometry of Einsteins's $*g$ -unified field theory*, International Journal of Theoretical Physics **27(9)** (1988).
- [6] K.T. Chung, *Conformal change in Einstein's $*g^{\lambda\nu}$ -unified field theory*, Nuove Cimento **(X)58B** (1968).
- [7] K.T. Chung, *Einstein's connection in terms of $*g^{\lambda\nu}$* , Nuove Cimento **(X) 27** (1963).
- [8] V. HLAVATÝ, *Geometry of Einstein's unified field theory*, P. Noordhoff Ltd (1957).

Department of Mathematics
Inha University
Incheon, 402-751, Korea
E-mail: `chcho@math.inha.ac.kr`