

## WEAK\* SMOOTH COMPACTNESS IN SMOOTH TOPOLOGICAL SPACES

CHUN-KEE PARK, WON KEUN MIN AND MYEONG HWAN KIM

ABSTRACT. In this paper we obtain some properties of the weak smooth  $\alpha$ -closure and weak smooth  $\alpha$ -interior of a fuzzy set in smooth topological spaces and introduce the concepts of several types of weak\* smooth compactness in smooth topological spaces and investigate some of their properties.

### 1. Introduction

Badard [1] introduced the concept of a smooth topological space which is a generalization of Chang's fuzzy topological space [2]. Many mathematical structures in smooth topological spaces were introduced and studied. In particular, Gayyar, Kerre and Ramadan [5] and Demirci [3, 4] introduced the concepts of smooth closure and smooth interior of a fuzzy set and several types of compactness in smooth topological spaces and obtained some of their properties. In [6] we introduced the concepts of smooth  $\alpha$ -closure and smooth  $\alpha$ -interior of a fuzzy set which are generalizations of smooth closure and smooth interior of a fuzzy set defined in [3] and also introduced several types of  $\alpha$ -compactness in smooth topological spaces and obtained some of their properties. In [7] we introduced the concepts of weak smooth  $\alpha$ -closure and weak smooth  $\alpha$ -interior of a fuzzy set and investigated some of their properties.

In this paper we obtain some properties of the weak smooth  $\alpha$ -closure and weak smooth  $\alpha$ -interior of a fuzzy set in smooth topological spaces and introduce the concepts of several types of weak\* smooth compactness in smooth topological spaces and investigate some of their properties.

---

Received July 7, 2003.

2000 Mathematics Subject Classification: 54A40, 03E72.

Key words and phrases: fuzzy sets, smooth topology, weak smooth  $\alpha$ -closure, weak smooth  $\alpha$ -interior, weak\* smooth compactness.

## 2. Preliminaries

Let  $X$  be a set and  $I = [0, 1]$  be the unit interval of the real line.  $I^X$  will denote the set of all fuzzy sets of  $X$ .  $0_X$  and  $1_X$  will denote the characteristic functions of  $\phi$  and  $X$ , respectively.

A smooth topological space (s.t.s.) [8] is an ordered pair  $(X, \tau)$ , where  $X$  is a non-empty set and  $\tau : I^X \rightarrow I$  is a mapping satisfying the following conditions:

$$(O1) \tau(0_X) = \tau(1_X) = 1;$$

$$(O2) \forall A, B \in I^X, \tau(A \cap B) \geq \tau(A) \wedge \tau(B);$$

$$(O3) \text{ for every subfamily } \{A_i : i \in J\} \subseteq I^X, \tau(\cup_{i \in J} A_i) \geq \wedge_{i \in J} \tau(A_i).$$

Then the mapping  $\tau : I^X \rightarrow I$  is called a smooth topology on  $X$ . The number  $\tau(A)$  is called the degree of openness of  $A$ .

A mapping  $\tau^* : I^X \rightarrow I$  is called a smooth cotopology [8] iff the following three conditions are satisfied:

$$(C1) \tau^*(0_X) = \tau^*(1_X) = 1;$$

$$(C2) \forall A, B \in I^X, \tau^*(A \cup B) \geq \tau^*(A) \wedge \tau^*(B);$$

$$(C3) \text{ for every subfamily } \{A_i : i \in J\} \subseteq I^X, \tau^*(\cap_{i \in J} A_i) \geq \wedge_{i \in J} \tau^*(A_i).$$

If  $\tau$  is a smooth topology on  $X$ , then the mapping  $\tau^* : I^X \rightarrow I$ , defined by  $\tau^*(A) = \tau(A^c)$  where  $A^c$  denotes the complement of  $A$ , is a smooth cotopology on  $X$ . Conversely, if  $\tau^*$  is a smooth cotopology on  $X$ , then the mapping  $\tau : I^X \rightarrow I$ , defined by  $\tau(A) = \tau^*(A^c)$ , is a smooth topology on  $X$  [8].

Demirci [3] introduced the concepts of smooth closure and smooth interior in smooth topological spaces as follows:

Let  $(X, \tau)$  be a s.t.s. and  $A \in I^X$ . Then the  $\tau$ -smooth closure (resp.,  $\tau$ -smooth interior) of  $A$ , denoted by  $\bar{A}$  (resp.,  $A^\circ$ ), is defined by  $\bar{A} = \cap\{K \in I^X : \tau^*(K) > 0, A \subseteq K\}$  (resp.,  $A^\circ = \cup\{K \in I^X : \tau(K) > 0, K \subseteq A\}$ ). Demirci [4] defined the families  $W(\tau) = \{A \in I^X : A = A^\circ\}$  and  $W^*(\tau) = \{A \in I^X : A = \bar{A}\}$ , where  $(X, \tau)$  is a s.t.s. Note that  $A \in W(\tau) \Leftrightarrow A^c \in W^*(\tau)$ .

Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces. A function  $f : X \rightarrow Y$  is called smooth continuous with respect to  $\tau$  and  $\sigma$  [8] iff  $\tau(f^{-1}(A)) \geq \sigma(A)$  for every  $A \in I^Y$ . A function  $f : X \rightarrow Y$  is called weakly smooth continuous with respect to  $\tau$  and  $\sigma$  [8] iff  $\sigma(A) > 0 \Rightarrow \tau(f^{-1}(A)) > 0$  for every  $A \in I^Y$ . In this paper, a weakly

smooth continuous function with respect to  $\tau$  and  $\sigma$  is called a quasi-smooth continuous function with respect to  $\tau$  and  $\sigma$ .

A function  $f : X \rightarrow Y$  is smooth continuous with respect to  $\tau$  and  $\sigma$  iff  $\tau^*(f^{-1}(A)) \geq \sigma^*(A)$  for every  $A \in I^Y$ . A function  $f : X \rightarrow Y$  is weakly smooth continuous with respect to  $\tau$  and  $\sigma$  iff  $\sigma^*(A) > 0 \Rightarrow \tau^*(f^{-1}(A)) > 0$  for every  $A \in I^Y$  [8].

A function  $f : X \rightarrow Y$  is called smooth open (resp., smooth closed) with respect to  $\tau$  and  $\sigma$  [8] iff  $\tau(A) \leq \sigma(f(A))$  (resp.,  $\tau^*(A) \leq \sigma^*(f(A))$ ) for every  $A \in I^X$ .

A function  $f : X \rightarrow Y$  is called smooth preserving (resp., strict smooth preserving) with respect to  $\tau$  and  $\sigma$  [5] iff  $\sigma(A) \geq \sigma(B) \Leftrightarrow \tau(f^{-1}(A)) \geq \tau(f^{-1}(B))$  (resp.,  $\sigma(A) > \sigma(B) \Leftrightarrow \tau(f^{-1}(A)) > \tau(f^{-1}(B))$ ) for every  $A, B \in I^Y$ .

If  $f : X \rightarrow Y$  is a smooth preserving function (resp., a strict smooth preserving function) with respect to  $\tau$  and  $\sigma$ , then  $\sigma^*(A) \geq \sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) \geq \tau^*(f^{-1}(B))$  (resp.,  $\sigma^*(A) > \sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) > \tau^*(f^{-1}(B))$ ) for every  $A, B \in I^Y$  [5].

A function  $f : X \rightarrow Y$  is called smooth open preserving (resp., strict smooth open preserving) with respect to  $\tau$  and  $\sigma$  [5] iff  $\tau(A) \geq \tau(B) \Rightarrow \sigma(f(A)) \geq \sigma(f(B))$  (resp.,  $\tau(A) > \tau(B) \Rightarrow \sigma(f(A)) > \sigma(f(B))$ ) for every  $A, B \in I^X$ .

Let  $(X, \tau)$  be a s.t.s.,  $\alpha \in [0, 1)$  and  $A \in I^X$ . The  $\tau$ -smooth  $\alpha$ -closure (resp.,  $\tau$ -smooth  $\alpha$ -interior) of  $A$ , denoted by  $\overline{A}_\alpha$  (resp.,  $A_\alpha^o$ ), is defined by  $\overline{A}_\alpha = \bigcap \{K \in I^X : \tau^*(K) > \alpha\tau^*(A), A \subseteq K\}$  (resp.,  $A_\alpha^o = \bigcup \{K \in I^X : \tau(K) > \alpha\tau(A), K \subseteq A\}$ ) [6]. In [7] we defined the families  $W_\alpha(\tau) = \{A \in I^X : A = A_\alpha^o\}$  and  $W_\alpha^*(\tau) = \{A \in I^X : A = \overline{A}_\alpha\}$ , where  $(X, \tau)$  is a s.t.s. Note that  $A \in W_\alpha(\tau) \Leftrightarrow A^c \in W_\alpha^*(\tau)$ .

### 3. weak smooth $\alpha$ -closure and weak smooth $\alpha$ -interior

In this section, we investigate some properties of the weak smooth  $\alpha$ -closure and weak smooth  $\alpha$ -interior of a fuzzy set in smooth topological spaces.

DEFINITION 3.1[7]. Let  $(X, \tau)$  be a s.t.s.,  $\alpha \in [0, 1)$  and  $A \in I^X$ . The weak  $\tau$ -smooth  $\alpha$ -closure (resp., weak  $\tau$ -smooth  $\alpha$ -interior) of  $A$ , denoted by  $wcl_\alpha(A)$  (resp.,  $wint_\alpha(A)$ ), is defined by  $wcl_\alpha(A) = \bigcap \{K \in I^X : K \in W_\alpha^*(\tau), A \subseteq K\}$  (resp.,  $wint_\alpha(A) = \bigcup \{K \in I^X : K \in$

$W_\alpha(\tau), K \subseteq A\}$ .

**THEOREM 3.2.** *Let  $(X, \tau)$  be a s.t.s.,  $\alpha \in [0, 1)$  and  $A \in I^X$ . Then*

- (a)  $A \subseteq wcl_\alpha(A) \subseteq \bar{A} \subseteq \bar{A}_\alpha$ ,
- (b)  $A_\alpha^o \subseteq A^o \subseteq wint_\alpha(A) \subseteq A$ .

*Proof.* (a) Let  $K \in I^X$  and  $A \subseteq K$ . Then  $\tau^*(K) > \alpha\tau^*(A) \Rightarrow \tau^*(K) > 0$  and  $\tau^*(K) > 0 \Rightarrow K = \bar{K}_\alpha$ , i.e.,  $K \in W_\alpha^*(\tau)$  by Theorem 3.6[6]. From the definitions of  $\bar{A}_\alpha$ ,  $\bar{A}$  and  $wcl_\alpha(A)$  we have  $A \subseteq wcl_\alpha(A) \subseteq \bar{A} \subseteq \bar{A}_\alpha$ .

(b) Let  $K \in I^X$  and  $K \subseteq A$ . Then  $\tau(K) > \alpha\tau(A) \Rightarrow \tau(K) > 0$  and  $\tau(K) > 0 \Rightarrow K = K_\alpha^o$ , i.e.,  $K \in W_\alpha(\tau)$  by Theorem 3.6[6]. From the definition of  $A_\alpha^o$ ,  $A^o$  and  $wint_\alpha(A)$  we have  $A_\alpha^o \subseteq A^o \subseteq wint_\alpha(A) \subseteq A$ .  $\square$

**THEOREM 3.3.** *Let  $(X, \tau)$  be a s.t.s.,  $\alpha \in [0, 1)$  and  $A, B \in I^X$ . Then*

- (a)  $A \subseteq B \Rightarrow wcl_\alpha(A) \subseteq wcl_\alpha(B)$ ,
- (b)  $A \subseteq B \Rightarrow wint_\alpha(A) \subseteq wint_\alpha(B)$ ,
- (c)  $(wcl_\alpha(A))^c = wint_\alpha(A^c)$ ,
- (d)  $wcl_\alpha(A) = (wint_\alpha(A^c))^c$ ,
- (e)  $(wint_\alpha(A))^c = wcl_\alpha(A^c)$ ,
- (f)  $wint_\alpha(A) = (wcl_\alpha(A^c))^c$ .

*Proof.* (a) and (b) follow directly from Definition 3.2.

(c) From Definition 3.2 we have

$$\begin{aligned} (wcl_\alpha(A))^c &= (\cap\{K \in I^X : K \in W_\alpha^*(\tau), A \subseteq K\})^c \\ &= \cup\{K^c : K \in I^X, K^c \in W_\alpha(\tau), K^c \subseteq A^c\} \\ &= \cup\{U \in I^X : U \in W_\alpha(\tau), U \subseteq A^c\} \\ &= wint_\alpha(A^c). \end{aligned}$$

(d), (e) and (f) can be easily obtained from (c).  $\square$

**DEFINITION 3.4[4].** Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces. A function  $f : X \rightarrow Y$  is called weak smooth continuous with respect to  $\tau$  and  $\sigma$  iff  $A \in W(\sigma) \Rightarrow f^{-1}(A) \in W(\tau)$  for every  $A \in I^Y$ .

Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces. A function  $f : X \rightarrow Y$  is weak smooth continuous with respect to  $\tau$  and  $\sigma$  iff  $A \in W^*(\sigma) \Rightarrow f^{-1}(A) \in W^*(\tau)$  for every  $A \in I^Y$  [4].

**THEOREM 3.5.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces. If a function  $f : X \rightarrow Y$  is quasi-smooth continuous with respect to  $\tau$  and  $\sigma$ , then  $f : X \rightarrow Y$  is weak smooth continuous with respect to  $\tau$  and  $\sigma$ .*

*Proof.* Let  $f : X \rightarrow Y$  be a quasi-smooth continuous function with respect to  $\tau$  and  $\sigma$ . Then by Proposition 3.5[3]  $f^{-1}(A^\circ) \subseteq (f^{-1}(A))^\circ$  for every  $A \in I^Y$ . Let  $A \in W(\sigma)$ , i.e.,  $A = A^\circ$ . Then  $f^{-1}(A) = f^{-1}(A^\circ) \subseteq (f^{-1}(A))^\circ$ . From the definition of smooth interior we have  $(f^{-1}(A))^\circ \subseteq f^{-1}(A)$ . Hence  $f^{-1}(A) = (f^{-1}(A))^\circ$ , i.e.,  $f^{-1}(A) \in W(\tau)$ . Therefore  $f : X \rightarrow Y$  is weak smooth continuous with respect to  $\tau$  and  $\sigma$ .  $\square$

**DEFINITION 3.6.** Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces and let  $\alpha \in [0, 1)$ . A function  $f : X \rightarrow Y$  is called weak smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$  iff  $A \in W_\alpha(\sigma) \Rightarrow f^{-1}(A) \in W_\alpha(\tau)$  for every  $A \in I^Y$ .

**THEOREM 3.7.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces and let  $\alpha \in [0, 1)$ . If a function  $f : X \rightarrow Y$  is weak smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$ , then*

- (a)  $f(wcl_\alpha(A)) \subseteq wcl_\alpha(f(A))$  for every  $A \in I^X$ ,
- (b)  $wcl_\alpha(f^{-1}(A)) \subseteq f^{-1}(wcl_\alpha(A))$  for every  $A \in I^Y$ ,
- (c)  $f^{-1}(wint_\alpha(A)) \subseteq wint_\alpha(f^{-1}(A))$  for every  $A \in I^Y$ .

*Proof.* (a) For every  $A \in I^X$ , we have

$$\begin{aligned}
 & f^{-1}(wcl_\alpha(f(A))) \\
 &= f^{-1}(\cap\{U \in I^Y : U \in W_\alpha^*(\sigma), f(A) \subseteq U\}) \\
 &\supseteq f^{-1}(\cap\{U \in I^Y : f^{-1}(U) \in W_\alpha^*(\tau), A \subseteq f^{-1}(U)\}) \\
 &= \cap\{f^{-1}(U) \in I^X : U \in I^Y, f^{-1}(U) \in W_\alpha^*(\tau), A \subseteq f^{-1}(U)\} \\
 &\supseteq \cap\{K \in I^X : K \in W_\alpha^*(\tau), A \subseteq K\} \\
 &= wcl_\alpha(A).
 \end{aligned}$$

Hence  $f(wcl_\alpha(A)) \subseteq wcl_\alpha(f(A))$ .

(b) For every  $A \in I^Y$ , we have

$$\begin{aligned}
& f^{-1}(wcl_\alpha(A)) \\
&= f^{-1}(\cap\{U \in I^Y : U \in W_\alpha^*(\sigma), A \subseteq U\}) \\
&\supseteq f^{-1}(\cap\{U \in I^Y : f^{-1}(U) \in W_\alpha^*(\tau), f^{-1}(A) \subseteq f^{-1}(U)\}) \\
&= \cap\{f^{-1}(U) \in I^X : U \in I^Y, f^{-1}(U) \in W_\alpha^*(\tau), \\
&\quad f^{-1}(A) \subseteq f^{-1}(U)\} \\
&\supseteq \cap\{K \in I^X : K \in W_\alpha^*(\tau), f^{-1}(A) \subseteq K\} \\
&= wcl_\alpha(f^{-1}(A)).
\end{aligned}$$

(c) For every  $A \in I^Y$ , we have

$$\begin{aligned}
& f^{-1}(wint_\alpha(A)) \\
&= f^{-1}(\cup\{U \in I^Y : U \in W_\alpha(\sigma), U \subseteq A\}) \\
&\subseteq f^{-1}(\cup\{U \in I^Y : f^{-1}(U) \in W_\alpha(\tau), f^{-1}(U) \subseteq f^{-1}(A)\}) \\
&= \cup\{f^{-1}(U) \in I^X : U \in I^Y, f^{-1}(U) \in W_\alpha(\tau), \\
&\quad f^{-1}(U) \subseteq f^{-1}(A)\} \\
&\subseteq \cup\{K \in I^X : K \in W_\alpha(\tau), K \subseteq f^{-1}(A)\} \\
&= wint_\alpha(f^{-1}(A)).
\end{aligned}$$

□

#### 4. Types of weak\* smooth compactness

In this section, we introduce the concepts of several types of weak\* smooth compactness in smooth topological spaces and investigate some of their properties.

We define the families  $W_{w\alpha}(\tau) = \{A \in I^X : A = wint_\alpha(A)\}$  and  $W_{w\alpha}^*(\tau) = \{A \in I^X : A = wcl_\alpha(A)\}$ , where  $(X, \tau)$  is a s.t.s. and  $\alpha \in [0, 1)$ . Then

$$\begin{aligned}
& A \in W_{w\alpha}(\tau) \Leftrightarrow A^c \in W_{w\alpha}^*(\tau), \\
& A \in W_\alpha(\tau) \Rightarrow A \in W(\tau) \Rightarrow A \in W_{w\alpha}(\tau), \\
& A \in W_\alpha^*(\tau) \Rightarrow A \in W^*(\tau) \Rightarrow A \in W_{w\alpha}^*(\tau).
\end{aligned}$$

DEFINITION 4.1. Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces and let  $\alpha \in [0, 1)$ . A function  $f : X \rightarrow Y$  is called weak\* smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$  iff  $A \in W_{w\alpha}(\sigma) \Rightarrow f^{-1}(A) \in W_{w\alpha}(\tau)$  for every  $A \in I^Y$ .

Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces. A function  $f : X \rightarrow Y$  is weak\* smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$  iff  $A \in W_{w\alpha}^*(\sigma) \Rightarrow f^{-1}(A) \in W_{w\alpha}^*(\tau)$  for every  $A \in I^Y$ .

DEFINITION 4.2. Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces and let  $\alpha \in [0, 1)$ . A function  $f : X \rightarrow Y$  is called weak\* smooth  $\alpha$ -open (resp., weak\* smooth  $\alpha$ -closed) with respect to  $\tau$  and  $\sigma$  iff  $A \in W_{w\alpha}(\tau) \Rightarrow f(A) \in W_{w\alpha}(\sigma)$  (resp.,  $A \in W_{w\alpha}^*(\tau) \Rightarrow f(A) \in W_{w\alpha}^*(\sigma)$ ) for every  $A \in I^X$ .

DEFINITION 4.3. Let  $\alpha \in [0, 1)$ . A s.t.s.  $(X, \tau)$  is called weak\* smooth compact iff every family in  $W_{w\alpha}(\tau)$  covering  $X$  has a finite subcover.

DEFINITION 4.4. Let  $\alpha \in [0, 1)$ . A s.t.s.  $(X, \tau)$  is called weak\* smooth nearly compact iff for every family  $\{A_i : i \in J\}$  in  $W_{w\alpha}(\tau)$  covering  $X$ , there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} \overline{A_i}^o = 1_X$ .

DEFINITION 4.5. Let  $\alpha \in [0, 1)$ . A s.t.s.  $(X, \tau)$  is called weak\* smooth almost compact iff for every family  $\{A_i : i \in J\}$  in  $W_{w\alpha}(\tau)$  covering  $X$ , there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} \overline{A_i} = 1_X$ .

THEOREM 4.6. Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces,  $\alpha \in [0, 1)$  and  $f : X \rightarrow Y$  a surjective and weak\* smooth  $\alpha$ -continuous function with respect to  $\tau$  and  $\sigma$ . If  $(X, \tau)$  is weak\* smooth compact, then so is  $(Y, \sigma)$ .

*Proof.* Let  $\{A_i : i \in J\}$  be a family in  $W_{w\alpha}(\sigma)$  covering  $Y$ , i.e.,  $\cup_{i \in J} A_i = 1_Y$ . Then  $\cup_{i \in J} f^{-1}(A_i) = f^{-1}(1_Y) = 1_X$ . Since  $f : X \rightarrow Y$  is weak\* smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$ ,  $\{f^{-1}(A_i) : i \in J\} \subseteq W_{w\alpha}(\tau)$ . Since  $(X, \tau)$  is weak\* smooth compact, there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} f^{-1}(A_i) = 1_X$ . From the surjectivity of  $f$  we have  $1_Y = f(1_X) = f(\cup_{i \in J_0} f^{-1}(A_i)) = \cup_{i \in J_0} f(f^{-1}(A_i)) = \cup_{i \in J_0} A_i$ . Therefore  $(Y, \sigma)$  is weak\* smooth compact.  $\square$

**THEOREM 4.7.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces and let  $\alpha \in [0, 1)$ . If a function  $f : X \rightarrow Y$  is weak smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$ , then  $f : X \rightarrow Y$  is weak\* smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$ .*

*Proof.* Let  $f : X \rightarrow Y$  be a weak smooth  $\alpha$ -continuous function with respect to  $\tau$  and  $\sigma$ . Then by Theorem 3.7  $f^{-1}(\text{wint}_\alpha(A)) \subseteq \text{wint}_\alpha(f^{-1}(A))$  for every  $A \in I^Y$ . Let  $A \in W_{w\alpha}(\sigma)$ , i.e.,  $A = \text{wint}_\alpha(A)$ . Then  $f^{-1}(A) = f^{-1}(\text{wint}_\alpha(A)) \subseteq \text{wint}_\alpha(f^{-1}(A))$ . From the definition of weak smooth  $\alpha$ -interior we have  $\text{wint}_\alpha(f^{-1}(A)) \subseteq f^{-1}(A)$ . Hence  $f^{-1}(A) = \text{wint}_\alpha(f^{-1}(A))$ , i.e.,  $f^{-1}(A) \in W_{w\alpha}(\tau)$ . Therefore  $f : X \rightarrow Y$  is weak\* smooth  $\alpha$ -continuous with respect to  $\tau$  and  $\sigma$ .  $\square$

We obtain the following corollary from Theorem 4.6 and 4.7.

**COROLLARY 4.8.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces,  $\alpha \in [0, 1)$  and  $f : X \rightarrow Y$  a surjective and weak smooth  $\alpha$ -continuous function with respect to  $\tau$  and  $\sigma$ . If  $(X, \tau)$  is weak\* smooth compact, then so is  $(Y, \sigma)$ .*

**THEOREM 4.9.** *Let  $\alpha \in [0, 1)$ . Then a weak\* smooth nearly compact s.t.s.  $(X, \tau)$  is weak\* smooth almost compact.*

*Proof.* Let  $\{A_i : i \in J\}$  be a family in  $W_{w\alpha}(\tau)$  covering  $X$ . Since  $(X, \tau)$  is weak\* smooth nearly compact, there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} (\overline{A_i})^o = 1_X$ . Since  $(\overline{A_i})^o \subseteq \overline{A_i}$  for each  $i \in J$  by Proposition 3.2[3],  $1_X = \cup_{i \in J_0} (\overline{A_i})^o \subseteq \cup_{i \in J_0} \overline{A_i}$ . So  $\cup_{i \in J_0} \overline{A_i} = 1_X$ . Hence  $(X, \tau)$  is weak\* smooth almost compact.  $\square$

**THEOREM 4.10.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces,  $\alpha \in [0, 1)$  and  $f : X \rightarrow Y$  a surjective and quasi-smooth continuous function with respect to  $\tau$  and  $\sigma$ . If  $(X, \tau)$  is weak\* smooth almost compact, then so is  $(Y, \sigma)$ .*

*Proof.* Let  $\{A_i : i \in J\}$  be a family in  $W_{w\alpha}(\sigma)$  covering  $Y$ , i.e.,  $\cup_{i \in J} A_i = 1_Y$ . Then  $1_X = f^{-1}(1_Y) = \cup_{i \in J} f^{-1}(A_i)$ . Since  $f$  is quasi-smooth continuous with respect to  $\tau$  and  $\sigma$ ,  $f$  is weak\* smooth continuous with respect to  $\tau$  and  $\sigma$  by Theorem 3.5 and 4.7. Hence  $f^{-1}(A_i) \in W_{w\alpha}(\tau)$  for each  $i \in J$ . Since  $(X, \tau)$  is weak\* smooth almost compact, there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} \overline{f^{-1}(A_i)} = 1_X$ .



From the surjectivity of  $f$  we have  $1_Y = f(1_X) = f(\overline{\cup_{i \in J_0} f^{-1}(A_i)}) = \cup_{i \in J_0} f(\overline{f^{-1}(A_i)})$ . Since  $f : X \rightarrow Y$  is quasi-smooth continuous with respect to  $\tau$  and  $\sigma$ , from Proposition 3.5[3] we have  $\overline{f^{-1}(A_i)} \subseteq f^{-1}(\overline{A_i})$  for each  $i \in J$ . Hence  $1_Y = \cup_{i \in J_0} f(\overline{f^{-1}(A_i)}) \subseteq \cup_{i \in J_0} f(f^{-1}(\overline{A_i})) = \cup_{i \in J_0} \overline{A_i}$ , i.e.,  $\cup_{i \in J_0} \overline{A_i} = 1_Y$ . Thus  $(Y, \sigma)$  is weak\* smooth almost compact.  $\square$

**THEOREM 4.11.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two smooth topological spaces,  $\alpha \in [0, 1)$  and  $f : X \rightarrow Y$  a surjective, quasi-smooth continuous and smooth open function with respect to  $\tau$  and  $\sigma$ . If  $(X, \tau)$  is weak\* smooth nearly compact, then so is  $(Y, \sigma)$ .*

*Proof.* Let  $\{A_i : i \in J\}$  be a family in  $W_{w\alpha}(\sigma)$  covering  $Y$ , i.e.,  $\cup_{i \in J} A_i = 1_Y$ . Then  $1_X = f^{-1}(1_Y) = \cup_{i \in J} f^{-1}(A_i)$ . Since  $f$  is quasi-smooth continuous with respect to  $\tau$  and  $\sigma$ ,  $f$  is weak\* smooth continuous with respect to  $\tau$  and  $\sigma$  by Theorem 3.5 and 4.7. Hence  $f^{-1}(A_i) \in W_{w\alpha}(\tau)$  for each  $i \in J$ . Since  $(X, \tau)$  is weak\* smooth nearly compact, there exists a finite subset  $J_0$  of  $J$  such that  $\cup_{i \in J_0} \overline{(f^{-1}(A_i))^o} = 1_X$ . From the surjectivity of  $f$  we have  $1_Y = f(1_X) = f(\cup_{i \in J_0} \overline{(f^{-1}(A_i))^o}) = \cup_{i \in J_0} f(\overline{(f^{-1}(A_i))^o})$ . Since  $f : X \rightarrow Y$  is smooth open with respect to  $\tau$  and  $\sigma$ , from Proposition 3.6[3] we have  $f(\overline{(f^{-1}(A_i))^o}) \subseteq \overline{(f(f^{-1}(A_i)))^o}$  for each  $i \in J$ . Since  $f : X \rightarrow Y$  is quasi-smooth continuous with respect to  $\tau$  and  $\sigma$ , from Proposition 3.5[3] we have  $\overline{f^{-1}(A_i)} \subseteq f^{-1}(\overline{A_i})$  for each  $i \in J$ . Hence  $1_Y = \cup_{i \in J_0} f(\overline{(f^{-1}(A_i))^o}) \subseteq \cup_{i \in J_0} \overline{(f(f^{-1}(A_i)))^o} \subseteq \cup_{i \in J_0} \overline{(f(f^{-1}(\overline{A_i})))^o} = \cup_{i \in J_0} \overline{A_i}^o$ , i.e.,  $\cup_{i \in J_0} \overline{A_i}^o = 1_Y$ . Thus  $(Y, \sigma)$  is weak\* smooth nearly compact.  $\square$

## References

1. R. Badard, *Smooth axiomatics*, First IFSA Congress, Palma de Mallorca (July 1986).
2. C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182-190.
3. M. Demirci, *On several types of compactness in smooth topological spaces*, Fuzzy Sets and Systems **90** (1997), 83-88.
4. ———, *Three topological structures of smooth topological spaces*, Fuzzy Sets and Systems **101** (1999), 185-190.
5. M. K. El Gayyar, E. E. Kerre and A. A. Ramadan, *Almost compactness and near compactness in smooth topological spaces*, Fuzzy Sets and Systems **62** (1994), 193-202.

6. C. K. Park, W. K. Min and M. H. Kim,  $\alpha$ -compactness in smooth topological spaces, *Int. J. Math. and Math. Sci.* (to appear).
7. ———, *Quasi-smooth  $\alpha$ -structure of smooth topological spaces*, (submitted).
8. A. A. Ramadan, *Smooth topological spaces*, *Fuzzy Sets and Systems* **48** (1992), 371-375.
9. L. A. Zadeh, *Fuzzy sets*, *Inform. and Control* **8** (1965), 338-353.

Department of Mathematics  
Kangwon National University  
Chuncheon 200-701, Korea  
*E-mail*: ckpark@kangwon.ac.kr

Department of Mathematics  
Kangwon National University  
Chuncheon 200-701, Korea  
*E-mail*: wkmin@cc.kangwon.ac.kr

Department of Mathematics  
Kangwon National University  
Chuncheon 200-701, Korea  
*E-mail*: kimmw@kangwon.ac.kr