

**THE EXISTENCE, NONEXISTENCE AND UNIQUENESS
OF GLOBAL POSITIVE COEXISTENCE OF A
NONLINEAR ELLIPTIC BIOLOGICAL INTERACTING
MODEL**

JOON HYUK KANG, JUNGHO LEE, AND YUN MYUNG OH

ABSTRACT. The purpose of this paper is to give a sufficient condition for the existence, nonexistence and uniqueness of coexistence of positive solutions to a rather general type of elliptic competition system of the Dirichlet problem on the bounded domain Ω in R^n . The techniques used in this paper are upper-lower solutions, maximum principles and spectrum estimates. The arguments also rely on some detailed properties for the solution of logistic equations. This result yields an algebraically computable criterion for the positive coexistence of competing species of animals in many biological models.

1. Introduction

The coexistence of steady states of competition interacting models with diffusion has been an object of intensive study in recent years. See, for example, lists of references in [1], [3], [4], [5], [9], [10], [14], [15], [14], [15]. The most general type of parabolic competition interacting system is

$$\begin{cases} u_t = \Delta u + ug(u, v), \\ v_t = \Delta v + vh(u, v), \end{cases}$$

where Δ is the Laplacian and u, v represent the densities of two competing species of animals. The terms Δu and Δv model dispersal by means of simple diffusion. We assume here that the C^1 functions g and h are relative growth rates satisfying the following so-called growth rate

Received February 28, 2004.

2000 Mathematics Subject Classification: 35A05, 35A07, 35B50, 35G30, 35J25 and 35K20.

Key words and phrases: competition model, coexistence state.

Research supported by Andrews University Faculty Research Grant 2000.

conditions:

- (G_1) $g_u(u, v) < 0, g_v(u, v) < 0, h_u(u, v) < 0, h_v(u, v) < 0,$
 (G_2) There exist constants $c_0 > 0, c_1 > 0$ such that $g(u, 0) \leq 0$ for $u \geq c_0$
 and $h(0, v) \leq 0$ for $u \geq c_1$.

The hypothesis (G_1) characterizes how the two species u and v interact with each other in terms of their relative growth rates. It is well known that condition (G_2) exhibits the so-called logistic pattern while the constants c_0 and c_1 are referred to as the carrying capacity.

The earlier literature on this line focused in the Neumann boundary value problem:

$$(1) \quad \begin{cases} u_t = \Delta u + ug(u, v), \\ v_t = \Delta v + vh(u, v), \\ \frac{\partial u(t, x)}{\partial n} = 0 = \frac{\partial v(t, x)}{\partial n} \text{ for } (t, x) \in [0, T] \times \partial\Omega, \end{cases}$$

and in its steady state, the elliptic system

$$(2) \quad \begin{cases} \Delta u + ug(u, v) = 0, \\ \Delta v + vh(u, v) = 0, \\ \frac{\partial u(x)}{\partial n} = 0 = \frac{\partial v(x)}{\partial n} \text{ for } x \in \partial\Omega, \end{cases}$$

where n denotes the unit out-normal along boundary $\partial\Omega$. The Neumann boundary conditions $\frac{\partial u(x)}{\partial n} = 0 = \frac{\partial v(x)}{\partial n}$ are interpreted as an assumption that both populations are staying inside, that there is no migratory flux across $\partial\Omega$. The goals of investigations along this line include finding out under what conditions on the nonlinearities g and h systems (1) and (2) have positive solutions $u > 0, v > 0$ and the possible uniqueness. Most of the work in this case were established by P. DeMottoni and F. Rothe in 1979 [7] and P. Brown in 1980 [2]. Their work in a large sense completes the avenue of investigation in the study of Neumann boundary value problems. Researchers thus have since turned their attention to the biologically and physically more important case that is the Dirichlet boundary condition:

$$(3) \quad \begin{cases} \Delta u + ug(u, v) = 0, \\ \Delta v + vh(u, v) = 0, \\ (u, v)|_{\partial\Omega} = (0, 0). \end{cases}$$

Biologically, this setting allows migration of these two populations across the boundary but they may not stay on $\partial\Omega$, where, for example $\partial\Omega$ is a

river. It was then found that the features known in the Neumann setting are not usually shared by those in the Dirichlet setting. The study in the latter setting, especially in the case of steady states like system (3), seems to be more difficult.

The goal of this paper is to answer the following questions about positive steady state to (3).

Problem 1 : What are the sufficient conditions for existence of steady state?

Problem 2 : Is it possible for either one of the species to be extinct?

Problem 3 : When is the coexistence state unique?

2. Preliminaries

In this section we state some preliminary results which will be useful for our later arguments.

DEFINITION 2.1. (*Upper and Lower solutions*)
The vector functions $(\bar{u}^1, \dots, \bar{u}^N), (\underline{u}^1, \dots, \underline{u}^N)$ form an upper/lower solution pair for the system

$$\begin{cases} \Delta u^i + g^i(u^1, \dots, u^N) = 0 & \text{in } \Omega \\ u^i = 0 & \text{on } \partial\Omega \end{cases}$$

if for $i = 1, \dots, N$

$$\begin{cases} \Delta \bar{u}^i + g^i(u^1, \dots, u^{i-1}, \bar{u}^i, u^{i+1}, \dots, u^N) \leq 0 \\ \Delta \underline{u}^i + g^i(u^1, \dots, u^{i-1}, \underline{u}^i, u^{i+1}, \dots, u^N) \geq 0 \\ \text{in } \Omega \text{ for } \underline{u}^j \leq u^j \leq \bar{u}^j, j \neq i, \end{cases}$$

and

$$\begin{aligned} \underline{u}^i &\leq \bar{u}^i & \text{on } \Omega \\ \underline{u}^i &\leq 0 \leq \bar{u}^i & \text{on } \partial\Omega. \end{aligned}$$

LEMMA 2.1. ([1])
If g^i in the Definition 2.1 are in C^1 and the system admits an upper/lower

solution pair $(\underline{u}^1, \dots, \underline{u}^N), (\bar{u}^1, \dots, \bar{u}^N)$, then there is a solution of the system in 2.1 with $\underline{u}^i \leq u^i \leq \bar{u}^i$ in Ω . If

$$\begin{aligned} \Delta \bar{u}^i + g^i(\bar{u}^1, \dots, \bar{u}^N) &\neq 0, \\ \Delta \underline{u}^i + g^i(\underline{u}^1, \dots, \underline{u}^N) &\neq 0 \end{aligned}$$

in Ω for $i = 1, \dots, N$, then $\underline{u}^i < u^i < \bar{u}^i$ in Ω .

LEMMA 2.2. (The first eigenvalue)([6])

$$(4) \quad \begin{cases} -\Delta u + q(x)u = \lambda u & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, \end{cases}$$

where $q(x)$ is a smooth function from Ω to R and Ω is a bounded domain in R^n .

(A) The first eigenvalue $\lambda_1(q)$ of (4), denoted by simply λ_1 when $q \equiv 0$, is simple with a positive eigenfunction.

(B) If $q_1(x) < q_2(x)$ for all $x \in \Omega$, then $\lambda_1(q_1) < \lambda_1(q_2)$.

(C) (Variational Characterization of the first eigenvalue)

$$\lambda_1(q) = \min_{\phi \in W_0^1(\Omega), \phi \neq 0} \frac{\int_{\Omega} (|\nabla \phi|^2 + q\phi^2) dx}{\int_{\Omega} \phi^2 dx}$$

We also need some information on the solutions of the following logistic equations.

LEMMA 2.3. ([14])

$$\begin{cases} \Delta u + uf(u) = 0 & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, u > 0, \end{cases}$$

where f is a decreasing C^1 function such that there exists $c_0 > 0$ such that $f(u) \leq 0$ for $u \geq c_0$ and Ω is a bounded domain in R^n .

(1) If $f(0) > \lambda_1$, then the above equation has a unique positive solution, where λ_1 is the first eigenvalue of $-\Delta$ with homogeneous boundary condition. We denote this unique positive solution as θ_f .

(2) If $f(0) \leq \lambda_1$, then the above equation does not have any positive solution.

3. Existence, Nonexistence and Uniqueness

We consider the system (3) with conditions (G_1) and (G_2) .

THEOREM 3.1. (A) *If $g(0, c_1) > \lambda_1$ and $h(c_0, 0) > \lambda_1$, then (3) has a solution (u, v) with*

$$\begin{aligned}\theta_{g(\cdot, c_1)} &< u < \theta_{g(\cdot, 0)} \\ \theta_{h(c_0, \cdot)} &< v < \theta_{h(0, \cdot)}.\end{aligned}$$

Conversely, any solution (u, v) of (3) with $u > 0, v > 0$ in Ω must satisfy these inequalities.

(B) *If $g(0, 0) \leq \lambda_1$ or $h(0, 0) \leq \lambda_1$, then (3) does not have any positive solution.*

Proof. (A) Let $\bar{u} = \theta_{g(\cdot, 0)}, \bar{v} = \theta_{h(0, \cdot)}$. Then by the monotonicity of g ,

$$\begin{aligned}\Delta \bar{u} + \bar{u}g(\bar{u}, \bar{v}) \\ &= \Delta \bar{u} + \bar{u}(g(\bar{u}, 0) - g(\bar{u}, 0) + g(\bar{u}, \bar{v})) \\ &= \bar{u}(g(\bar{u}, \bar{v}) - g(\bar{u}, 0)) < 0.\end{aligned}$$

Similarly,

$$\Delta \bar{v} + \bar{v}h(\bar{u}, \bar{v}) < 0.$$

So, (\bar{u}, \bar{v}) is an upper solution to (3).

Let $\underline{u} = \theta_{g(\cdot, c_1)}$ and $\underline{v} = \theta_{h(c_0, \cdot)}$. Then by the Maximum Principles, we obtain

$$\begin{aligned}\underline{u} &\leq \theta_{g(\cdot, 0)} \leq c_0, \\ \underline{v} &\leq \theta_{h(0, \cdot)} \leq c_1.\end{aligned}$$

By the monotonicity of g ,

$$\begin{aligned}\Delta \underline{u} + \underline{u}g(\underline{u}, \underline{v}) \\ &= \Delta \underline{u} + \underline{u}(g(\underline{u}, c_1) - g(\underline{u}, c_1) + g(\underline{u}, \underline{v})) \\ &= \underline{u}(g(\underline{u}, \underline{v}) - g(\underline{u}, c_1)) \geq 0.\end{aligned}$$

Similarly,

$$\Delta \underline{v} + \underline{v}h(\underline{u}, \underline{v}) \geq 0.$$

Therefore, $(\underline{u}, \underline{v})$ is a lower solution to (3). Furthermore, $\underline{u} < \bar{u}, \underline{v} < \bar{v}$ in Ω and $\underline{u} = \bar{u} = \underline{v} = \bar{v} = 0$ on $\partial\Omega$.

So, (3) has a solution (u, v) with

$$\begin{aligned}\theta_{g(\cdot, c_1)} &< u < \theta_{g(\cdot, 0)}, \\ \theta_{h(c_0, \cdot)} &< v < \theta_{h(0, \cdot)}.\end{aligned}$$

Suppose (u, v) is a coexistence state for (3). Then since

$$\begin{aligned} & \Delta u + ug(u, 0) \\ \geq & \Delta u + ug(u, v) = 0, \end{aligned}$$

u is a lower solution of

$$(5) \quad \begin{aligned} \Delta Z + Zg(Z, 0) &= 0 \text{ in } \Omega, \\ Z &= 0 \text{ on } \partial\Omega. \end{aligned}$$

But, since any constant larger than c_0 is an upper solution of (5), we have

$$(6) \quad u < \theta_{g(\cdot, 0)}.$$

Similarly, we have

$$(7) \quad v < \theta_{h(0, \cdot)}.$$

Since $v < \theta_{h(0, \cdot)} \leq c_1$, by the monotonicity of g

$$\begin{aligned} & \Delta u + ug(u, c_1) \\ \leq & \Delta u + ug(u, v) = 0. \end{aligned}$$

Therefore, u is an upper solution of

$$(8) \quad \begin{aligned} \Delta Z + Zg(Z, c_1) &= 0 \text{ in } \Omega, \\ Z &= 0 \text{ on } \partial\Omega. \end{aligned}$$

If $\epsilon > 0$ is so small that $g(\epsilon\phi_1, c_1) > \lambda_1$ on $\bar{\Omega}$, where ϕ_1 is the first eigenvector of $-\Delta$ with homogeneous boundary condition, then since

$$\begin{aligned} & \Delta\epsilon\phi_1 + \epsilon\phi_1g(\epsilon\phi_1, c_1) \\ &= \epsilon(\Delta\phi_1 + \phi_1g(\epsilon\phi_1, c_1)) \\ &> \epsilon(\Delta\phi_1 + \lambda_1\phi_1) = 0, \end{aligned}$$

$\epsilon\phi_1$ is a lower solution of (8). So, we have

$$(9) \quad \theta_{g(\cdot, c_1)} < u.$$

Similarly, we have

$$(10) \quad \theta_{h(c_0, \cdot)} < v.$$

By (6), (7), (9) and (10),

$$\begin{aligned} \theta_{g(\cdot, c_1)} &< u < \theta_{g(\cdot, 0)}, \\ \theta_{h(c_0, \cdot)} &< v < \theta_{h(0, \cdot)}. \end{aligned}$$

(B) Assume $g(0, 0) \leq \lambda_1$. The other cases are proved similarly. Suppose (\bar{u}, \bar{v}) is a positive solution to (3). Then since

$$\begin{aligned} & \Delta \bar{u} + \bar{u}g(\bar{u}, 0) \\ &= \Delta \bar{u} + \bar{u}(g(\bar{u}, \bar{v}) - g(\bar{u}, \bar{v}) + g(\bar{u}, 0)) \\ &= \bar{u}(g(\bar{u}, 0) - g(\bar{u}, \bar{v})) \geq 0, \end{aligned}$$

\bar{u} is a lower solution to

$$(11) \quad \begin{aligned} \Delta u + ug(u, 0) &= 0 \text{ in } \Omega, \\ u &= 0 \text{ on } \partial\Omega. \end{aligned}$$

Any constant larger than c_0 is an upper solution to (11). Hence, (11) has a positive solution u_0 with $\bar{u} < u_0$. This contradicts to the Lemma 2.3 which says there is no positive solution of (11) if $g(0, 0) \leq \lambda_1$. \square

THEOREM 3.2. *If $g(0, c_1) > \lambda_1, h(c_0, 0) > \lambda_1$ and*

$$4 \inf\left(-\frac{\partial g(u, v)}{\partial u}\right) \inf\left(-\frac{\partial h(u, v)}{\partial v}\right) \geq \frac{\theta_{g(\cdot, 0)}}{\theta_{h(c_0, \cdot)}} \left(\sup \frac{\partial g(u, v)}{\partial v}\right)^2 + \frac{\theta_{h(0, \cdot)}}{\theta_{g(\cdot, c_1)}} \left(\sup \frac{\partial h(u, v)}{\partial u}\right)^2 + 2\left(\sup \frac{\partial g(u, v)}{\partial v}\right)\left(\sup \frac{\partial h(u, v)}{\partial u}\right),$$

then (3) has a unique positive solution.

Proof. Suppose (u_1, v_1) and (u_2, v_2) are positive solutions to (3). Let $p = u_1 - u_2$ and $q = v_1 - v_2$. Then

$$\begin{aligned} & \Delta p + pg(u_1, v_1) \\ &= \Delta u_1 - \Delta u_2 + (u_1 - u_2)g(u_1, v_1) \\ &= -\Delta u_2 - u_2g(u_1, v_1) \\ &= -\Delta u_2 - u_2(g(u_2, v_2) - g(u_2, v_2) + g(u_1, v_1)) \\ &= -u_2(g(u_1, v_1) - g(u_2, v_2)) \\ &= -u_2(g(u_1, v_1) - g(u_2, v_1) + g(u_2, v_1) - g(u_2, v_2)). \end{aligned}$$

But, by the Mean Value Theorem, there is \tilde{x} depending on u_1, u_2 such that

$$g(u_1, v_1) - g(u_2, v_1) = \frac{\partial g(\tilde{x}, v_1)}{\partial u} p.$$

Hence,

$$\Delta p + pg(u_1, v_1) = -u_2 \left[\frac{\partial g(\tilde{x}, v_1)}{\partial u} p + g(u_2, v_1) - g(u_2, v_2) \right].$$

i.e.,

$$(12) \quad \Delta p + g(u_1, v_1)p + u_2 p \frac{\partial g(\tilde{x}, v_1)}{\partial u} - u_2(g(u_2, v_2) - g(u_2, v_1)) = 0.$$

The same argument shows that

$$(13) \quad \Delta q + h(u_2, v_2)q + v_1 q \frac{\partial h(u_2, \bar{x})}{\partial v} - v_1(h(u_2, v_1) - h(u_1, v_1)) = 0,$$

where \bar{x} depends on v_1, v_2 by the Mean Value Theorem.

Since $\lambda_1(-g(u_1, v_1)) = 0$, by the Variational Characterization of the first eigenvalue,

$$(14) \quad \int_{\Omega} Z(-\Delta Z - g(u_1, v_1)Z)dx \geq 0$$

for any $Z \in C^2(\bar{\Omega})$ and $Z|_{\partial\Omega} = 0$. The same argument shows that

$$(15) \quad \int_{\Omega} W(-\Delta W - h(u_2, v_2)W)dx \geq 0$$

for any $W \in C^2(\bar{\Omega})$ and $W|_{\partial\Omega} = 0$.

From (12) and (13), we get

$$\begin{aligned} -p\Delta p - g(u_1, v_1)p^2 - \frac{\partial g(\bar{x}, v_1)}{\partial u} u_2 p^2 + u_2 p(g(u_2, v_2) - g(u_2, v_1)) &= 0, \\ -q\Delta q - h(u_2, v_2)q^2 - \frac{\partial h(u_2, \bar{x})}{\partial v} v_1 q^2 + v_1 q(h(u_2, v_1) - h(u_1, v_1)) &= 0. \end{aligned}$$

Hence from (14) and (15),

$$\int_{\Omega} \left(-\frac{\partial g(\bar{x}, v_1)}{\partial u} u_2 p^2 + u_2 p(g(u_2, v_2) - g(u_2, v_1)) + v_1 q(h(u_2, v_1) - h(u_1, v_1)) - \frac{\partial h(u_2, \bar{x})}{\partial v} v_1 q^2 \right) dx \leq 0.$$

By the Mean Value Theorem, for each $x \in \Omega$, there exist \tilde{y}, \bar{y} such that

$$\begin{aligned} g(u_2, v_2) - g(u_2, v_1) &= \frac{\partial g(u_2, \tilde{y})}{\partial v} (-q), \\ h(u_2, v_1) - h(u_1, v_1) &= \frac{\partial h(\bar{y}, v_1)}{\partial u} (-p), \end{aligned}$$

which implies that

$$\int_{\Omega} -\frac{\partial g(\tilde{x}, v_1)}{\partial u} u_2 p^2 - \left(u_2 \frac{\partial g(u_2, \tilde{y})}{\partial v} + v_1 \frac{\partial h(\bar{y}, v_1)}{\partial u} \right) pq - \frac{\partial h(u_2, \bar{x})}{\partial v} v_1 q^2 dx \leq 0.$$

Therefore, we find

$$\begin{aligned} p \equiv q \equiv 0 \quad \text{if} \quad & -\frac{\partial g(\tilde{x}, v_1)}{\partial u} u_2 \zeta^2 - \left(u_2 \frac{\partial g(u_2, \tilde{y})}{\partial v} + v_1 \frac{\partial h(\bar{y}, v_1)}{\partial u} \right) \zeta \eta \\ & - \frac{\partial h(u_2, \bar{x})}{\partial v} v_1 \eta^2 \quad \text{is positive definite} \\ & \text{for each } x \in \Omega. \end{aligned}$$

This is the case if

$$\begin{aligned} & u_2^2 \left(\frac{\partial g(u_2, \tilde{y})}{\partial v} \right)^2 + v_1^2 \left(\frac{\partial h(\bar{y}, v_1)}{\partial u} \right)^2 + 2u_2 v_1 \frac{\partial g(u_2, \tilde{y})}{\partial v} \frac{\partial h(\bar{y}, v_1)}{\partial u} \\ & - 4 \frac{\partial g(\tilde{x}, v_1)}{\partial u} \frac{\partial h(u_2, \bar{x})}{\partial v} u_2 v_1 \leq 0 \quad \text{for each } x \in \Omega. \end{aligned}$$

$$\text{i.e., } 4 \frac{\partial g(\bar{x}, v_1)}{\partial u} \frac{\partial h(u_2, \bar{x})}{\partial v} \geq \frac{u_2}{v_1} \left(\frac{\partial g(u_2, \bar{y})}{\partial v} \right)^2 + \frac{v_1}{u_2} \left(\frac{\partial h(\bar{y}, v_1)}{\partial u} \right)^2 + 2 \frac{\partial g(u_2, \bar{y})}{\partial v} \frac{\partial h(\bar{y}, v_1)}{\partial u} \text{ for each } x \in \Omega.$$

But, from the inequality in (A) and the hypothesis in the theorem,

$$\begin{aligned} \frac{u_2}{v_1} \left(\frac{\partial g(u_2, \bar{y})}{\partial v} \right)^2 &+ 2 \frac{\partial g(u_2, \bar{y})}{\partial v} \frac{\partial h(\bar{y}, v_1)}{\partial u} + \frac{v_1}{u_2} \left(\frac{\partial h(\bar{y}, v_1)}{\partial u} \right)^2 \\ &\leq \frac{\theta_{g(\cdot, 0)}}{\theta_{h(c_0, \cdot)}} \left(\sup \frac{\partial g(u, v)}{\partial v} \right)^2 + \frac{\theta_{h(0, \cdot)}}{\theta_{g(\cdot, c_1)}} \left(\sup \frac{\partial h(u, v)}{\partial u} \right)^2 \\ &\quad + 2 \sup \left(\frac{\partial g(u, v)}{\partial v} \right) \sup \left(\frac{\partial h(u, v)}{\partial u} \right) \\ &\leq 4 \inf \left(- \frac{\partial g(u, v)}{\partial u} \right) \inf \left(- \frac{\partial h(u, v)}{\partial v} \right) \\ &\leq 4 \frac{\partial g(\bar{x}, v_1)}{\partial u} \frac{\partial g(u_2, \bar{x})}{\partial v}. \end{aligned}$$

□

We can also extend the results to the case when there are multiple species competing in the same environment.

Consider the interacting model

$$(16) \quad \begin{aligned} \Delta u_i + u_i g_i(u_1, u_2, \dots, u_i, u_{i+1}, \dots, u_N) &= 0 \text{ in } \Omega, \\ u_i &= 0 \text{ on } \partial\Omega \end{aligned}$$

for $i = 1, \dots, N$.

Again, we assume here that the C^1 functions g_i for $i = 1, \dots, N$ are relative growth rates satisfying the following growth rate conditions:

$$(M1) \quad \frac{\partial g_i}{\partial u_j} < 0 \text{ for } i, j = 1, 2, \dots, N,$$

(M2) There exist constants $c_1 > 0, c_2 > 0, \dots, c_N > 0$ such that $g_i(0, \dots, 0, u_i, 0, \dots, 0) \leq 0$ for $u_i \geq c_i$.

Again, (M1) characterizes how the N species u_1, u_2, \dots, u_N interact with each other in terms of their relative growth rates and (M2) is the logistic pattern with carrying capacity constants c_1, c_2, \dots, c_N .

The followings are the main results. The proofs are similar to those with 2 competing species, and so we just sketch it without the details.

THEOREM 3.3. (A) *If $g_i(c_1, c_2, \dots, c_{i-1}, 0, c_{i+1}, \dots, c_N) > \lambda_1$ for $i = 1, \dots, N$, then (16) has a solution (u_1, \dots, u_N) with*

$$\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)} < u_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$$

for $i = 1, \dots, N$.

Conversely, any solution (u_1, \dots, u_N) of (16) with $u_i > 0$ in Ω must satisfy these inequalities.

(B) If $g_i(0, \dots, 0) \leq \lambda_1$ for some $i = 1, \dots, N$, then (16) does not have any positive solution.

Proof. (A) Let $\bar{u}_i = \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$ and $\underline{u}_i = \theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)}$ for $i = 1, \dots, N$. Then by the Maximum Principles and the monotonicity of g_i , $(\bar{u}_1, \dots, \bar{u}_i, \dots, \bar{u}_N)$ and $(\underline{u}_1, \dots, \underline{u}_i, \dots, \underline{u}_N)$ are upper and lower solutions to (16), respectively.

Furthermore, for $i = 1, \dots, N$, $\underline{u}_i < \bar{u}_i$ in Ω and $\underline{u}_i = \bar{u}_i = 0$ on $\partial\Omega$. So, (16) has a solution (u_1, \dots, u_N) with the desired inequalities

$$\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)} < u_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$$

for $i = 1, \dots, N$.

Suppose (u_1, \dots, u_N) is a coexistence state for (16). Then by the direct computation using the monotonicity of g_i , we know that u_i is a lower solution of

$$(17) \quad \begin{aligned} \Delta Z + Zg_i(0, \dots, 0, Z, 0, \dots, 0) &= 0 \text{ in } \Omega, \\ Z &= 0 \text{ on } \partial\Omega \end{aligned}$$

for $i = 1, \dots, N$

But, since any constant larger than c_i is an upper solution of (17), we have

$$(18) \quad u_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$$

for $i = 1, \dots, N$.

Since $u_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)} \leq c_i$, by the monotonicity of g_i , we can derive that u_i is an upper solution of

$$(19) \quad \begin{aligned} \Delta Z + Zg_i(c_1, \dots, c_{i-1}, Z, c_{i+1}, \dots, c_N) &= 0 \text{ in } \Omega, \\ Z &= 0 \text{ on } \partial\Omega \end{aligned}$$

for $i = 1, \dots, N$.

If $\epsilon > 0$ is so small that $g_i(c_1, \dots, c_{i-1}, \epsilon\phi_1, c_{i+1}, \dots, c_N) > \lambda_1$ on $\bar{\Omega}$, where ϕ_1 is the first eigenvector of $-\Delta$ with homogeneous boundary condition, then by the direct computation again, we know that $\epsilon\phi_1$ is a lower solution of (19).

So, we have

$$(20) \quad \theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)} < u_i$$

for $i = 1, \dots, N$.

By (18) and (20),

$$\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)} < u_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$$

for $i = 1, \dots, N$.

(B) Without loss of generality, assume $g_1(0, \dots, 0) \leq \lambda_1$.

Suppose $(\bar{u}_1, \dots, \bar{u}_N)$ is a positive solution to (16). Then by the monotonicity of g_i , \bar{u}_1 is a lower solution to

$$(21) \quad \begin{aligned} \Delta Z + Zg_1(Z, 0, \dots, 0) &= 0 \text{ in } \Omega, \\ Z &= 0 \text{ on } \partial\Omega. \end{aligned}$$

Hence, by the fact that any constant larger than c_1 is an upper solution to (21), (21) has a positive solution u_1 with $\bar{u}_1 < u_1$ that contradicts to the Lemma 2.3. \square

THEOREM 3.4. *If $g_i(c_1, \dots, c_{i-1}, 0, c_{i+1}, \dots, c_N) > \lambda_1$ and*

$$2 \inf\left(-\frac{\partial g_i}{\partial x_i}\right) > \sum_{j=1, j \neq i}^N \left(\sup\left(-\frac{\partial g_i}{\partial x_j}\right) + K \sup\left(-\frac{\partial g_j}{\partial x_i}\right)\right)$$

for $i = 1, \dots, N$, where $K = \sup_{i, j \neq i} \frac{\theta_{g_j(0, \dots, 0, \dots, 0, \dots, 0)}}{\theta_{g_i(c_1, \dots, c_{i-1}, \dots, c_{i+1}, \dots, c_N)}}$, then (16) has a unique coexistence state.

Proof. Suppose (u_1, \dots, u_N) and (v_1, \dots, v_N) are coexistence states of (16) and let $w_i = u_i - v_i$ for $i = 1, \dots, N$. Then by the direct computation and the Variational Characterization of the first eigenvalue, we obtain

$$\int_{\Omega} \sum_{i=1}^N [v_i w_i (g_i(v_1, \dots, v_i, \dots, v_N) - g_i(u_1, \dots, u_i, \dots, u_N))] dx \leq 0.$$

By the Mean Value Theorem, there exist t^i and z^{ij} such that

$$(22) \quad \int_{\Omega} \sum_{i=1}^N \left[\frac{\partial g_i(v_1, \dots, v_{i-1}, t^i, v_{i+1}, \dots, v_N)}{\partial x_i} (-v_i) w_i^2 + \sum_{j=1, j \neq i}^N v_i w_i \frac{\partial g_i(u_1, \dots, u_{j-1}, z^{ij}, v_{j+1}, \dots, v_N)}{\partial x_j} (-w_j) \right] dx \leq 0.$$

If the integrand in the left side of (22) is positive definite, then (22) implies that $w_i \equiv 0$ in Ω for $i = 1, \dots, N$, which means the uniqueness of the coexistence state for (16). But for any $\epsilon > 0$,

$$\begin{aligned} & \frac{\partial g_i(u_1, \dots, u_{j-1}, z^{ij}, v_{j+1}, \dots, v_N)}{\partial x_j} (-v_i) w_i(w_j) \\ & \leq \frac{\partial g_i(u_1, \dots, u_{j-1}, z^{ij}, v_{j+1}, \dots, v_N)}{\partial x_j} (-v_i) \left[\frac{w_i^2}{2\epsilon} + \frac{\epsilon w_j^2}{2} \right]. \end{aligned}$$

So, we can see that the integrand is positive definite if for $i = 1, \dots, N$ and $x \in \Omega$,

$$> \sum_{j=1, j \neq i}^N \left(\frac{\frac{\partial g_i(v_1, \dots, v_{i-1}, t^i, v_{i+1}, \dots, v_N)}{\partial x_j}(-v_i)}{2\epsilon} + \frac{\epsilon \frac{\partial}{\partial x_i} g_j(u_1, \dots, u_{i-1}, z^{j^i}, v_{i+1}, \dots, v_N)(-v_j)}{2} \right)$$

or equivalently,

$$(23) \quad > \sum_{j=1, j \neq i}^N \left(\frac{-\frac{\partial g_i(v_1, \dots, v_{i-1}, t^i, v_{i+1}, \dots, v_N)}{\partial x_j} g_i(u_1, \dots, u_{j-1}, z^{ij}, v_{j+1}, \dots, v_N)}{2\epsilon} - \frac{\epsilon \frac{\partial}{\partial x_i} g_j(u_1, \dots, u_{i-1}, z^{ji}, v_{i+1}, \dots, v_N) \frac{v_j}{v_i}}{2} \right).$$

Since $\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)} < v_i < \theta_{g_i(0, \dots, 0, \cdot, 0, \dots, 0)}$ in Ω for $i = 1, \dots, N$, (23) will hold if for $i = 1, \dots, N$,

$$> \sum_{j=1, j \neq i}^N \left(\frac{-\frac{\partial g_i(v_1, \dots, v_{i-1}, t^i, v_{i+1}, \dots, v_N)}{\partial x_j}}{2\epsilon} + \frac{\epsilon \sup(-\frac{\partial g_j}{\partial x_i})}{2} \frac{\theta_{g_j(0, \dots, 0, \cdot, 0, \dots, 0)}}{\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)}} \right).$$

Let $K = \sup_{i, j \neq i} \frac{\theta_{g_j(0, \dots, 0, \cdot, 0, \dots, 0)}}{\theta_{g_i(c_1, \dots, c_{i-1}, \cdot, c_{i+1}, \dots, c_N)}}$. Then (23) holds if

$$\inf(-\frac{\partial g_i}{\partial x_i}) > \sum_{j=1, j \neq i}^N \left(\frac{\sup(-\frac{\partial g_j}{\partial x_j})}{2\epsilon} + \frac{K \epsilon \sup(-\frac{\partial g_j}{\partial x_i})}{2} \right).$$

Choosing $\epsilon = 1$, we have

$$2 \inf(-\frac{\partial g_i}{\partial x_i}) > \sum_{j=1, j \neq i}^N \left(\sup(-\frac{\partial g_j}{\partial x_j}) + K \sup(-\frac{\partial g_j}{\partial x_i}) \right).$$

□

References

- [1] S. W. Ali and C. Cosner, On the uniqueness of the positive steady state for Lotka - Volterra Models with diffusion, *Journal of Mathematical Analysis and Application* 168, 329-341(1992)
- [2] P. Brown, Decay to uniform states in ecological interactions, *SIAM J. Appl. Math.* 38:22-37(1980)
- [3] R. S. Cantrell and C. Cosner, On the steady - state problem for the Volterra - Lotka competition model with diffusion, *Houston Journal of mathematics*, 13(1987), 337-352.

- [4] R. S. Cantrell and C. Cosner, On the uniqueness and stability of positive solutions in the Volterra-Lotka competition model with diffusion, *Houston J. Math.* 15(1989) 341-361.
- [5] C. Cosner and A. C. Lazer, Stable coexistence states in the Volterra-Lotka competition model with diffusion, *Siam J. Appl. Math.*, 44(1984), 1112-1132.
- [6] D. Dunninger, Lecture note for applied analysis in Michigan State University
- [7] P. DeMottoni and F. Rothe, Convergence to homogenous equilibrium states for generalized Volterra-Lotka systems, *SIAM J. Appl. Math.* 648-663(1979)
- [8] R. Courant and D. Hilbert, *Methods of mathematical physics, Vol.1*, Interscience, New York, 1961.
- [9] C. Gui and Y. Lou, Uniqueness and nonuniqueness of coexistence states in the Lotka-Volterra competition model, *Comm Pure and Appl. Math.*, Vol.XVL2, No. 12(1994), 1571-1594.
- [10] J. L. Gomez and J. P. Pardo, Existence and uniqueness for some competition models with diffusion, *C.R. Acad. Sci. Paris*, 313 Série 1(1991), 933-938.
- [11] P. Hess, On uniqueness of positive solutions of nonlinear elliptic boundary value problems, *Math. Z.*, 165(1977), 17-18.
- [12] Joon H. Kang, Some global interacting system(submitted)
- [13] Joon H. Kang and Yun M. Oh, A sufficient condition for the uniqueness of positive steady state to a reaction diffusion system(to be appeared in the *Journal of the Korean Mathematical Society*)
- [14] L. Li and R. Logan, Positive solutions to general elliptic competition models, *Differential and Integral Equations*, 4(1991), 817-834.
- [15] A. Leung, Equilibria and stabilities for competing-species, reaction-diffusion equations with Dirichlet boundary data, *J. Math. Anal. Appl.*, 73(1980), 204-218.
- [16] M. H. Protter and H.F. Weinberger, *Maximum principles in differential equations*, Prentice Hall, Englewood Cliffs, N. J., 1967.
- [17] I. Stakgold and L. E. Payne, *Nonlinear problems in nuclear reactor analysis, in nonlinear problems in the physical sciences and biology, Lecture notes in Mathematics 322*, Springer, Berlin, 1973, 298-307.

Joon Hyuk Kang
Department of Mathematics
Andrews University
Berrien Springs, MI 49104
U.S.A.
E-mail: kang@andrews.edu

Jungho Lee
Department of Mathematics
Michigan State University
East Lansing, MI 48823
U.S.A.

Yun Myung Oh
Department of Mathematics
Michigan State University
East Lansing, MI 48823
U.S.A.
E-mail: ohy@math.msu.edu